CERN-PH-TH/2013-047
KA-TP-06-2013
PSI-PR-13-04

arXiv:1303.3876v2 [hep-ph] 26 Mar 2013

Effective Lagrangian for a light Higgs-like scalar

Roberto Contino[a], Margherita Ghezzi[a], Christophe Grojean[b,c],
Margarete Mühlleitner[d] and Michael Spira[e]

[a] *Dipartimento di Fisica, Università di Roma La Sapienza and INFN, Roma, Italy*
[b] *ICREA at IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Spain*
[c] *Theory Division, Physics Department, CERN, Geneva, Switzerland*
[d] *Institute for Theoretical Physics, Karlsruhe Institute of Technology, Karlsruhe, Germany*
[e] *Paul Scherrer Institut, CH–5232 Villigen PSI, Switzerland*

Abstract

We reconsider the effective Lagrangian that describes a light Higgs-like boson and better clarify a few issues which were not exhaustively addressed in the previous literature. In particular we highlight the strategy to determine whether the dynamics responsible for the electroweak symmetry breaking is weakly or strongly interacting. We also discuss how the effective Lagrangian can be implemented into automatic tools for the calculation of Higgs decay rates and production cross sections. Finally, we present eHDECAY, a modified version of the program HDECAY which includes the full list of leading bosonic operators of the effective Lagrangian and implements two benchmark composite Higgs models.

1 Introduction

The exploration of the weak scale has marked an important step forward with the discovery by the ATLAS [1] and CMS [2] collaborations of a boson with mass $m_h \simeq 125\,\mathrm{GeV}$, whose production cross section and decay rates are compatible with those predicted for the Higgs boson of the Standard Model (SM). At the same time, no hint of the existence of additional new particles has emerged yet, which might shed light on the origin of the electroweak symmetry breaking (EWSB). One is thus faced with the problem of which is the best strategy to describe the properties and investigate the nature of the new boson h, beyond the framework of the Standard Model. In absence of a direct observation of new states, our ignorance of the EWSB sector can be parametrized in terms of an effective Lagrangian for the light boson. Such an effective description is valid as long as New Physics (NP) states appear at a scale $M \gg m_h$, and is based on an expansion in the number of fields and derivatives [3]. The detailed form of the effective Lagrangian depends on which assumptions are made. Considering that the observation made by the LHC experiments is in remarkable agreement with the SM prediction, although within the current limited experimental precision, it is reasonable to assume that h is a CP-even scalar that forms an $SU(2)_L$ doublet together with the longitudinal polarizations of the W and Z, so that the $SU(2)_L \times U(1)_Y$ electroweak symmetry is linearly realized at high energies. Under these assumptions the effective Lagrangian can be expanded into a sum of operators with increasing dimensionality, where the leading NP effects are given by dimension-6 operators.

We reconsider such an effective Lagrangian in Section 2. By means of a naive power counting we estimate the coefficients of the various operators and review the most important bounds set on them by present experimental results on electroweak (EW) and flavor observables. Focusing on Higgs physics, we then discuss in Section 3 the relative effect of the various operators on physical observables. Such an analysis, first proposed in Ref. [4], will allow us to identify which operators can probe the Higgs coupling strength to the new states and which instead are sensitive only to the mass scale M. This is of key importance to distinguish between weakly-coupled UV completions of the Standard Model, like Supersymmetric (SUSY) theories, and theories where the EW symmetry is broken by a new strongly-interacting dynamics which forms the Higgs boson as a bound state [5–7, 4]. These are the two most compelling scenarios put forward to solve the hierarchy problem of the Standard Model. We conclude the section by discussing how the assumption of a Higgs doublet and linearly-realized $SU(2)_L \times U(1)_Y$ can be relaxed. We illustrate the non-linear effective Lagrangian valid for the case of a generic CP-even scalar h and discuss the implications of custodial invariance. Section 4 is devoted to clarify a few issues related to the use of the effective Lagrangian beyond the tree level. We present our concluding discussion in Section 5. In the Appendices A-C we collect useful formulas and give further details on the construction of the effective Lagrangian. The details of how we derived the bounds on the dimension-6 operators are reported in Appendix D. As an illustration of our analysis and to better demonstrate how the effective Lagrangian can be implemented into automatic tools for the computation of physical quantities like Higgs production cross sections and decay rates, in Appendix E we present `eHDECAY` [1], a modified version of the program `HDECAY` which includes the full list of leading bosonic operators.

2 Effective Lagrangian for a light Higgs doublet

The most general $SU(3)_C \times SU(2)_L \times U(1)_Y$-invariant Lagrangian for a weak doublet H at the level of dimension-6 operators was first classified in a systematic way in Refs. [8]. Subsequent analyses [9, 10] pointed out the presence of some redundant operators, and a minimal and complete list of operators was finally provided in Ref. [11]. As recently discussed in Ref. [4], a convenient basis of operators relevant for Higgs physics is the following:

$$\mathcal{L} = \mathcal{L}_{SM} + \sum_i \bar{c}_i O_i \equiv \mathcal{L}_{SM} + \Delta\mathcal{L}_{SILH} + \Delta\mathcal{L}_{F_1} + \Delta\mathcal{L}_{F_2} \tag{2.1}$$

---

[1] `eHDECAY` is available at the following URL: `http://www-itp.particle.uni-karlsruhe.de/~maggie/eHDECAY/`

with

$$\begin{aligned}
\Delta\mathcal{L}_{SILH} =\ & \frac{\bar{c}_H}{2v^2}\,\partial^\mu\big(H^\dagger H\big)\,\partial_\mu\big(H^\dagger H\big) + \frac{\bar{c}_T}{2v^2}\left(H^\dagger \overleftrightarrow{D^\mu} H\right)\left(H^\dagger \overleftrightarrow{D}_\mu H\right) - \frac{\bar{c}_6\,\lambda}{v^2}\left(H^\dagger H\right)^3 \\
& + \left(\frac{\bar{c}_u}{v^2}\,y_u\,H^\dagger H\,\bar{q}_L H^c u_R + \frac{\bar{c}_d}{v^2}\,y_d\,H^\dagger H\,\bar{q}_L H d_R + \frac{\bar{c}_l}{v^2}\,y_l\,H^\dagger H\,\bar{L}_L H l_R + h.c.\right) \\
& + \frac{i\bar{c}_W\,g}{2m_W^2}\left(H^\dagger\sigma^i\overleftrightarrow{D^\mu}H\right)(D^\nu W_{\mu\nu})^i + \frac{i\bar{c}_B\,g'}{2m_W^2}\left(H^\dagger\overleftrightarrow{D^\mu}H\right)(\partial^\nu B_{\mu\nu}) \\
& + \frac{i\bar{c}_{HW}\,g}{m_W^2}\,(D^\mu H)^\dagger\sigma^i(D^\nu H)W^i_{\mu\nu} + \frac{i\bar{c}_{HB}\,g'}{m_W^2}\,(D^\mu H)^\dagger(D^\nu H)B_{\mu\nu} \\
& + \frac{\bar{c}_\gamma\,{g'}^2}{m_W^2}\,H^\dagger H B_{\mu\nu}B^{\mu\nu} + \frac{\bar{c}_g\,g_S^2}{m_W^2}\,H^\dagger H G^a_{\mu\nu}G^{a\mu\nu}\,,
\end{aligned} \tag{2.2}$$

$$\begin{aligned}
\Delta\mathcal{L}_{F_1} =\ & \frac{i\bar{c}_{Hq}}{v^2}\,(\bar{q}_L\gamma^\mu q_L)\left(H^\dagger\overleftrightarrow{D}_\mu H\right) + \frac{i\bar{c}'_{Hq}}{v^2}\left(\bar{q}_L\gamma^\mu\sigma^i q_L\right)\left(H^\dagger\sigma^i\overleftrightarrow{D}_\mu H\right) \\
& + \frac{i\bar{c}_{Hu}}{v^2}\,(\bar{u}_R\gamma^\mu u_R)\left(H^\dagger\overleftrightarrow{D}_\mu H\right) + \frac{i\bar{c}_{Hd}}{v^2}\left(\bar{d}_R\gamma^\mu d_R\right)\left(H^\dagger\overleftrightarrow{D}_\mu H\right) \\
& + \left(\frac{i\bar{c}_{Hud}}{v^2}\,(\bar{u}_R\gamma^\mu d_R)\left(H^{c\,\dagger}\overleftrightarrow{D}_\mu H\right) + h.c.\right) \\
& + \frac{i\bar{c}_{HL}}{v^2}\left(\bar{L}_L\gamma^\mu L_L\right)\left(H^\dagger\overleftrightarrow{D}_\mu H\right) + \frac{i\bar{c}'_{HL}}{v^2}\left(\bar{L}_L\gamma^\mu\sigma^i L_L\right)\left(H^\dagger\sigma^i\overleftrightarrow{D}_\mu H\right) \\
& + \frac{i\bar{c}_{Hl}}{v^2}\left(\bar{l}_R\gamma^\mu l_R\right)\left(H^\dagger\overleftrightarrow{D}_\mu H\right),
\end{aligned} \tag{2.3}$$

$$\begin{aligned}
\Delta\mathcal{L}_{F_2} =\ & \frac{\bar{c}_{uB}\,g'}{m_W^2}\,y_u\,\bar{q}_L H^c\sigma^{\mu\nu}u_R\,B_{\mu\nu} + \frac{\bar{c}_{uW}\,g}{m_W^2}\,y_u\,\bar{q}_L\sigma^i H^c\sigma^{\mu\nu}u_R\,W^i_{\mu\nu} + \frac{\bar{c}_{uG}\,g_S}{m_W^2}\,y_u\,\bar{q}_L H^c\sigma^{\mu\nu}\lambda^a u_R\,G^a_{\mu\nu} \\
& + \frac{\bar{c}_{dB}\,g'}{m_W^2}\,y_d\,\bar{q}_L H\sigma^{\mu\nu}d_R\,B_{\mu\nu} + \frac{\bar{c}_{dW}\,g}{m_W^2}\,y_d\,\bar{q}_L\sigma^i H\sigma^{\mu\nu}d_R\,W^i_{\mu\nu} + \frac{\bar{c}_{dG}\,g_S}{m_W^2}\,y_d\,\bar{q}_L H\sigma^{\mu\nu}\lambda^a d_R\,G^a_{\mu\nu} \\
& + \frac{\bar{c}_{lB}\,g'}{m_W^2}\,y_l\,\bar{L}_L H\sigma^{\mu\nu}l_R\,B_{\mu\nu} + \frac{\bar{c}_{lW}\,g}{m_W^2}\,y_l\,\bar{L}_L\sigma^i H\sigma^{\mu\nu}l_R\,W^i_{\mu\nu} + h.c.
\end{aligned} \tag{2.4}$$

The SM Lagrangian $\mathcal{L}_{SM}$ and our convention for the covariant derivatives and the gauge field strengths are reported for completeness in Appendix A. In particular, λ is the Higgs quartic coupling and the weak scale is defined to be

$$v \equiv \frac{1}{(\sqrt{2}G_F)^{1/2}} = 246\,\mathrm{GeV}\,. \tag{2.5}$$

By $iH^\dagger \overleftrightarrow{D^\mu} H$ we denote the Hermitian derivative $iH^\dagger(D^\mu H) - i(D^\mu H)^\dagger H$, and $\sigma^{\mu\nu} \equiv i[\gamma^\mu, \gamma^\nu]/2$. The Yukawa couplings $y_{u,d,l}$ and the Wilson coefficients $\bar{c}_i$ in Eq. (2.3) are matrices in flavor space, and a sum over flavors has been left understood. As specified in Eq. (2.1), we will denote as O_i the dimension-6 operator whose coefficient is proportional to $\bar{c}_i$.

Our higher-dimensional Lagrangian, which is supposed to capture the leading New Physics effects, counts $12\,(\Delta\mathcal{L}_{SILH}) + 8\,(\Delta\mathcal{L}_{F_1}) + 8\,(\Delta\mathcal{L}_{F_2}) = 28$ operators. Five extra bosonic operators,

$$\begin{gathered}
\frac{\bar{c}_{3W}\, g^3}{m_W^2}\, \epsilon^{ijk} W_\mu^{i\,\nu} W_\nu^{j\,\rho} W_\rho^{k\,\mu} \,, \qquad \frac{\bar{c}_{3G}\, g_S^3}{m_W^2}\, f^{abc} G_\mu^{a\,\nu} G_\nu^{b\,\rho} G_\rho^{c\,\mu} \,, \\
\frac{\bar{c}_{2W}}{m_W^2}\, (D^\mu W_{\mu\nu})^i\, (D_\rho W^{\rho\nu})^i \;, \qquad \frac{\bar{c}_{2B}}{m_W^2}\, (\partial^\mu B_{\mu\nu})\, (\partial_\rho B^{\rho\nu}) \;, \qquad \frac{\bar{c}_{2G}}{m_W^2}\, (D^\mu G_{\mu\nu})^a\, (D_\rho G^{\rho\nu})^a \;,
\end{gathered} \tag{2.6}$$

which affect the gauge-boson propagators and self-interactions but with no effect on Higgs physics, should also be added to complete the operator basis, as well as 22 four-Fermi baryon-number-conserving operators. [2] A comparison with Ref. [11] shows that two of our operators are actually redundant. As we shall explain in more detail in Section 3 (see Eqs. (3.28), (3.29)), it is well known [10, 13] that two particular linear combinations of the fermionic operators in $\Delta\mathcal{L}_{F_1}$ are equivalent to pure oblique corrections parametrized by the operators O_T, O_W and O_B:

$$O_{H\Psi}^Y \equiv \sum_\psi Y_\psi\, O_{H\psi} \sim O_T\,, O_B \qquad \text{and} \qquad O'_{Hq} + O'_{HL} \sim O_W\,, \tag{2.7}$$

where the sum runs over all fermion representations, $\psi = q_L, u_R, d_R, L_L, l_R$, whose hypercharge has been denoted as Y_ψ. These two linear combinations have then to be excluded from $\Delta\mathcal{L}_{F_1}$, and we end up with exactly 53 linearly-independent operators as in Ref. [11]. [3] Any other dimension-six operator can be obtained from these 53 operators by using the equations of motion, or equivalently by performing appropriate field redefinitions. [4]

---

[2] Notice that the last three operators in Eq. (2.6) can be rewritten in favor of three additional independent four-Fermi operators, as in the basis of Ref. [11]. The coefficients $\bar{c}_{2W}$, $\bar{c}_{2B}$ contribute respectively to the W and Y parameters defined in Ref. [12].

[3] For completeness we collect in Appendix C also the extra 6 bosonic operators of dimension-six that are CP-odd.

Even though our basis (2.2)–(2.4) is equivalent to the one proposed in Ref. [11], we advocate that it is more appropriate for Higgs physics for at least three reasons [4]: *i)* Generic models of New Physics generate a contribution to the oblique $\hat{S}$ parameter [14, 12] at tree-level, which in the basis of Ref. [11] would have to be encoded in the two fermionic operators $O^Y_{H\psi}$ and $O'_{Hq} + O'_{HL}$ even in the absence of direct couplings between the SM fermions and the New Physics sector. There is an advantage in describing the oblique corrections in terms of the operators in (2.2) rather than in terms of the operators with fermionic currents, which generate vertex corrections and modify the Fermi constant. *ii)* The basis (2.1) isolates the contributions to the decays $h \to \gamma\gamma$ (from O_γ) and $h \to \gamma Z$ (from O_γ and $O_{HW} - O_{HB}$) that occur only at the radiative level in minimally coupled theories. *iii)* Our basis of operators is more appropriate to establish the nature of the Higgs boson and determine the strength of its interactions. For example, as we shall explain momentarily, if the Higgs boson is a pseudo Nambu-Goldstone boson (pNGB) the coefficient of the operator O_γ, hence the rate $h \to \gamma\gamma$, is suppressed, while in the basis of Ref. [11] this reflects into a cancellation in the linear combination $4\bar{c}_\gamma + (\bar{c}_{WW} - \bar{c}_{WB})$ (cf. footnote 4).

While a complete classification of the operators is essential, having a power counting to estimate their impact on physical observables, hence their relative importance, is equally crucial. In this sense a simple yet consequential observation was made in Ref. [4]: when expanding the effective Lagrangian in the number of fields and derivatives, any additional power of H is suppressed by a factor $g_*/M \equiv 1/f$, where $g_* \leq 4\pi$ denotes the coupling strength of the Higgs boson to New Physics states and M is their overall mass scale; any additional derivative instead costs a factor $1/M$. If the light Higgs boson is a composite state of the dynamics at the scale M, it is natural to expect $g_* \gg 1$, hence $f \ll M$, which implies that operators with extra powers of H give the leading corrections to low-energy

---

[4] In particular, the following identities hold:

$$
\begin{aligned}
&\frac{g^2}{4m_W^2}\, H^\dagger H\, W^i_{\mu\nu} W^{i\,\mu\nu} \equiv O_{WW} = O_W - O_B + O_{HB} - O_{HW} + \frac{1}{4}\, O_\gamma \\
&\frac{gg'}{4m_W^2}\, H^\dagger \sigma^i H\, W^i_{\mu\nu} B^{\mu\nu} \equiv O_{WB} = O_B - O_{HB} - \frac{1}{4}\, O_\gamma\,.
\end{aligned}
\tag{2.8}
$$

observables. On the other hand, in weakly-coupled completions of the Standard Model where $g_* \sim g$, all operators with the same dimension can be equally important. A proper analysis of the experimental results through the language of the effective Lagrangian can thus give indication on whether the dynamics at the origin of electroweak symmetry breaking is weakly or strongly interacting. According to the power counting of Ref. [4], one naively estimates $(\psi = u, d, l, q, L)$ [5]

$$\begin{aligned}
&\bar{c}_H, \bar{c}_T, \bar{c}_6, \bar{c}_\psi \sim O\left(\frac{v^2}{f^2}\right), \quad \bar{c}_W, \bar{c}_B \sim O\left(\frac{m_W^2}{M^2}\right), \quad \bar{c}_{HW}, \bar{c}_{HB}, \bar{c}_\gamma, \bar{c}_g \sim O\left(\frac{m_W^2}{16\pi^2 f^2}\right) \\
&\bar{c}_{H\psi}, \bar{c}'_{H\psi} \sim O\left(\frac{\lambda_\psi^2}{g_*^2}\frac{v^2}{f^2}\right), \quad \bar{c}_{Hud} \sim O\left(\frac{\lambda_u \lambda_d}{g_*^2}\frac{v^2}{f^2}\right), \quad \bar{c}_{\psi W}, \bar{c}_{\psi B}, \bar{c}_{\psi G} \sim O\left(\frac{m_W^2}{16\pi^2 f^2}\right),
\end{aligned} \tag{2.9}$$

where λ_ψ denotes the coupling of a generic SM fermion ψ to the new dynamics. It should be stressed that these estimates are valid at the UV scale M, at which the effective Lagrangian is matched onto explicit models. Renormalization effects between M and the EW scale mix operators with the same quantum numbers, and give in general subdominant corrections to the coefficients. We shall comment on these renormalization effects in Section 4. Notice that the estimates of $\bar{c}_{W,B}$, $\bar{c}_{H\psi}$, $\bar{c}'_{H\psi}$ and $\bar{c}_T$ apply when these coefficients are generated at tree-level. However, specific symmetry protections which might be at work in the UV theory, like for example R-parity in SUSY theories, can force the leading corrections to arise at the 1-loop level.

Equation (2.9) suggests that in the case of a strongly-interacting light Higgs boson (SILH) the leading New Physics effects in Higgs observables are parametrized by the operators $O_{H,T,6,\psi}$, and, if the SM fermions couple strongly to the new dynamics, by the fermionic operators of Eq. (2.3) [4]. Notice that, compared to the naive counting, $\bar{c}_{HW,HB,g,\gamma}$ are suppressed by an additional factor $(g_*^2/16\pi^2)$. This is because the corresponding operators contribute to the coupling of on-shell photons and gluons to neutral particles and modify the gyromagnetic ratio of the W, and are thus generated only at the loop level in a minimally coupled theory. Similarly, the dipole operators of Eq. (2.4) are generated at the loop-level only, hence their estimates have an extra loop factor.

---

[5] Notice that our normalization differs from the one of Ref. [4], and it is more convenient than the latter for a model-independent implementation of Eq. (2.2) in a computer program. The factor multiplying each operator in the effective Lagrangian has been conveniently defined such that the dependence on M and g_* is fully encoded in the dimensionless coefficients $\bar{c}_i$.

A special and phenomenologically motivated case is represented by theories where the Higgs doublet is a composite Nambu–Goldstone (NG) boson of a spontaneously-broken symmetry $\mathcal{G} \to \mathcal{H}$ of the strong dynamics [5–7,4]. For these models the scale f must be identified with the decay constant associated with the spontaneous breaking, and the naive estimate of the Wilson coefficients $\bar{c}_i$ is modified by the request of invariance under $\mathcal{G}$ in the limit of vanishing explicit breaking. At the level of dimension-6 operators, O_γ, O_g, O_6, $O_{u,d,l}$ and the dipole operators of Eq. (2.4) violate the shift symmetry $H^i \to H^i + \zeta^i$ ($\zeta^i = const.$) that is included as part of the $\mathcal{G}/\mathcal{H}$ transformations. This means that they cannot be generated in absence of an explicit breaking of the global symmetry. It follows, in particular, that the naive estimates of the operators O_γ and O_g carry in this case an additional suppression factor [4],

$$\bar{c}_\gamma, \bar{c}_g \sim O\left(\frac{m_W^2}{16\pi^2 f^2}\right) \times \frac{g_{\not{G}}^2}{g_*^2} \,, \tag{2.10}$$

where $g_{\not{G}}$ denotes any weak coupling that breaks the Goldstone symmetry (one of the SM weak couplings in minimal models, i.e. the SM gauge couplings or the Yukawa couplings). The operators O_6, O_ψ, $O_{\psi G}$, $O_{\psi W}$, $O_{\psi B}$ have been defined so that their prefactor already includes one spurion coupling, precisely the Higgs quartic coupling λ in O_6, and the Yukawa coupling y_ψ in the other operators – indeed, both these couplings vanish for an exact NG boson. The estimates of the corresponding coefficients $\bar{c}_6$, $\bar{c}_\psi$, $\bar{c}_{\psi G}$, $\bar{c}_{\psi W}$, $\bar{c}_{\psi B}$ are thus not modified.

In writing Eq. (2.2) we have assumed that each of the operators $O_{u,d,l}$ is flavor-aligned with the corresponding fermion mass term, as required in order to avoid large Flavor-Changing Neutral Currents (FCNC) mediated by the tree-level exchange of the Higgs boson (see for example Ref. [15] for a natural way to obtain this alignment). This implies one coefficient for the up-type quarks ($\bar{c}_u$), one for down-type quarks ($\bar{c}_d$), and one for the charged leptons ($\bar{c}_l$), i.e. the $\bar{c}_{u,d,l}$ are proportional to the identity matrix in flavor space.

2.1 Current bounds on flavor-preserving operators

It is useful to review some of the most important constraints on the coefficients $\bar{c}_i$ that follow from current experimental results, such as electroweak precision tests, flavor data and low-energy precision measurements. For simplicity, we focus on the bounds on flavor-conserving operators, keeping in mind that they can come also from flavor-changing processes. For a discussion of the bounds on flavor-violating operators see for example the recent review of Ref. [16] as well as Ref. [17].

Among the strongest bounds are those on operators that modify the vector-boson self-energies. The operator O_T, for example, violates the custodial symmetry [18] and contributes to the EW parameter ϵ_1 [19]. From the EW fit performed in Ref. [20], it follows, with 95% probability,

$$\Delta\epsilon_1 \equiv \Delta\rho = \bar{c}_T(m_Z)\,, \qquad -1.5\times 10^{-3} < \bar{c}_T(m_Z) < 2.2\times 10^{-3}\,. \tag{2.11}$$

Such a stringent bound can be more naturally satisfied by assuming that the dynamics at the scale M possesses an (at least approximate) $SU(2)_V$ custodial invariance. In this case $c_T(M)=0$, and a non-vanishing value will be generated through the renormalization-group (RG) flow of this Wilson coefficient down to m_Z in the presence of an explicit breaking of the custodial symmetry, as due for example to the Yukawa or hypercharge couplings. We will discuss these renormalization effects in more detail in Section 4. Notice that all the other dimension-6 operators in the effective Lagrangian are (formally) custodially symmetric and their coefficients will not be suppressed at the scale M. [6] The electroweak precision tests also imply a strong bound on $O_W + O_B$ [4], since this linear combination contributes to the parameter ϵ_3 [19]. With 95% probability, one has [20]:

$$\Delta\epsilon_3 = \bar{c}_W(m_Z) + \bar{c}_B(m_Z)\,, \qquad -1.4\times 10^{-3} < \bar{c}_W(m_Z) + \bar{c}_B(m_Z) < 1.9\times 10^{-3}\,. \tag{2.12}$$

From the tree-level estimate of $\bar{c}_{W,B}$ reported in Eq. (2.9), and assuming an approximate custodial invariance to suppress $\bar{c}_T$ as explained above, it follows that Eqs. (2.11) and (2.12) set a lower bound $M \gtrsim$ a few TeV. This bound is quite robust and can be avoided only in weakly-coupled UV completions where an extra symmetry protection suppresses the leading contribution to $\bar{c}_{W,B}$ by an additional loop factor. Notable examples are SUSY theories with R-parity.

---

[6] More precisely, for all the other operators the only violation of the custodial symmetry comes from the explicit breaking due to the gauging of hypercharge. As such, this breaking is external to the EWSB dynamics, since it comes from the weak gauging of its global symmetries. Formal invariance of the operators can be restored by uplifting the hypercharge gauge field to a whole triplet of $SU(2)_R$. The top Yukawa coupling is another source of explicit custodial breaking.

The fermionic operators in Eq. (2.3) are strongly constrained by Z-pole measurements, as they modify the couplings of the Z to quarks and leptons:

$$\frac{\delta g_{L\psi}}{g_{L\psi}} = \frac{1}{2}\,\frac{\bar{c}_{H\Psi} + 2\,T_{3L}\,\bar{c}'_{H\Psi}}{T_{3L} - Q\sin^2\theta_W}\,, \qquad \frac{\delta g_{R\psi}}{g_{R\psi}} = \frac{1}{2}\,\frac{\bar{c}_{H\psi}}{Q\sin^2\theta_W}\,, \tag{2.13}$$

where T_{3L} and Q are respectively the $SU(2)_L$ and electric charges of the fermion ψ, and $\Psi = \{L, q\}$ is the $SU(2)_L$ doublet to which ψ_L belongs. We used the results of Ref. [20] to perform a fit on the coefficients $\bar{c}_{H\psi}$, $\bar{c}_{H\Psi}$, $\bar{c}'_{H\Psi}$. The details of our analysis can be found in Appendix D (see also Ref. [21]). In the case of light quarks (u, d, s) we find the following bounds

$$\begin{gathered}
-0.03 < \bar{c}_{Hq1} < 0.02\,, \quad -0.002 < \bar{c}'_{Hq1} < 0.003\,, \\
-0.005 < \bar{c}_{Hq2} < 0.003\,, \quad -0.003 < \bar{c}'_{Hq2} < 0.005\,, \\
-0.008 < \bar{c}_{Hu} < 0.02\,, \quad -0.03 < \bar{c}_{Hd} < 0.02\,, \quad -0.03 < \bar{c}_{Hs} < 0.02\,,
\end{gathered} \tag{2.14}$$

while a fit on leptons and heavy quarks (c, b) gives

$$\begin{gathered}
-0.004 < \bar{c}_{HL} + \bar{c}'_{HL} < 0.002\,, \quad -0.003 < \bar{c}_{HL} - \bar{c}'_{HL} < 0.0002\,, \quad -0.0007 < \bar{c}_{Hl} < 0.003\,, \\
-0.02 < \bar{c}_{Hq_2} + \bar{c}'_{Hq_2} < 0.005\,, \quad -0.02 < \bar{c}_{Hc} < 0.03\,, \\
-0.003 < \bar{c}_{Hq_3} - \bar{c}'_{Hq_3} < 0.009\,, \quad -0.07 < \bar{c}_{Hb} < -0.005\,.
\end{gathered} \tag{2.15}$$

All the above bounds have 95% probability and by the various coefficients we mean their values at the scale m_Z. The weakest constraint is that on the operator O_{Hb}, which modifies the coupling of b_R to the Z boson. The operator involving two right-handed top quarks, O_{Ht}, is unconstrained by EW data, but it is also not relevant for the Higgs decays and will be neglected in the following. The coefficient $\bar{c}_{Htb}$ is severely constrained by the $b \to s\gamma$ rate. Indeed, the expansion of O_{Htb} around the vacuum contains a vertex of the type $W t_R b_R$, which at 1-loop gives a chirally-enhanced contribution to the rate (see for example Ref. [22]). We find, with 95% probability:

$$-\,0.4 \times 10^{-3} < \bar{c}_{Htb}(m_W) < 1.3 \times 10^{-3}\,. \tag{2.16}$$

For a given (v/f), the above bounds set a limit on the couplings of the SM fermions to the new dynamics, see Eq. (2.9). Unless the scale of New Physics is very large, or some specific symmetry protection is at work in the UV theory (see for example the discussion in Ref. [21]), it follows that the SM fermions must be very weakly coupled to the new dynamics, with the exception of the top quark.

The constraints on the dipole operators of Eq. (2.4) come from the current experimental limits on electric dipole moments (EDMs) and anomalous magnetic moments. The bounds on the neutron and mercury EDMs for example strongly constrain the dipole operators with u and d quarks. By using the formulas of Ref. [23] we find, with 95% probability, that:

$$\begin{aligned} -7.01\times 10^{-6} < \mathrm{Im}(\bar{c}_{uB}+\bar{c}_{uW}) < 7.86\times 10^{-6}\,, \\ -9.42\times 10^{-7} < \mathrm{Im}(\bar{c}_{dB}-\bar{c}_{dW}) < 8.40\times 10^{-7}\,, \\ -1.62\times 10^{-6} < \mathrm{Im}(\bar{c}_{uG}) < 2.01\times 10^{-6}\,, \\ -7.71\times 10^{-7} < \mathrm{Im}(\bar{c}_{dG}) < 5.70\times 10^{-7}\,, \end{aligned} \tag{2.17}$$

where the coefficients are evaluated at the low-energy scale $\mu \sim 1\,\mathrm{GeV}$. According to the naive estimate (2.9), for $O(1)$ CP-violating phases these results imply a bound on $(v/f)^2$ at the level of 10^{-3}. In natural extensions of the SM, such a strong limit clearly points to the need of a symmetry protection mechanism. For a discussion, see for example Ref. [21] for the case of composite Higgs theories, and Ref. [24] for the case of SUSY theories.

Among the heavier quarks the most interesting bounds are those on dipole operators with top quarks [25]. These come from the experimental limit on the neutron EDM,

$$-1.39\times 10^{-4} < \mathrm{Im}(\bar{c}_{tG}) < 1.21\times 10^{-4}\,, \tag{2.18}$$

the $b\to s\gamma$ and $b\to s l^+ l^-$ rates,

$$-0.057 < \mathrm{Re}(\bar{c}_{tW}+\bar{c}_{tB}) - 2.65\,\mathrm{Im}(\bar{c}_{tW}+\bar{c}_{tB}) < 0.20\,, \tag{2.19}$$

and the $t\bar{t}$ cross sections measured at the Tevatron and LHC,

$$-6.12\times 10^{-3} < \mathrm{Re}(\bar{c}_{tG}) < 1.94\times 10^{-3}\,. \tag{2.20}$$

All these bounds have 95% probability and have been derived by making use of the formulas reported in Ref. [25]. [7] It is worth noting that the bounds of Eqs. (2.19) and (2.20) are still about one order of magnitude weaker than the size of $\bar{c}_{tG}$, $\bar{c}_{tW}$ and $\bar{c}_{tB}$ expected from the naive estimate (2.9) with $(v/f)^2 \sim 0.1$. Additional weaker constraints arise from the limits on anomalous top interactions based on top decays and single top production. From the results of Ref. [26] we find that, with 95% probability:

$$-1.2 < \mathrm{Re}(\bar{c}_{bW}) < 1.1\,, \qquad -0.01 < \mathrm{Re}(\bar{c}_{tW}) < 0.02\,. \tag{2.21}$$

where the coefficients are evaluated at the scale $\mu \sim m_t$.

In the lepton sector, the current measurements and SM predictions of the muon [27, 28] and electron [29, 30] anomalous magnetic moments and the limits on their EDMs [31, 32] imply the following 95% probability bounds:

$$\begin{gathered} -1.64 \times 10^{-2} < \mathrm{Re}(\bar{c}_{eB} - \bar{c}_{eW}) < 3.37 \times 10^{-3}\,, \\ 1.88 \times 10^{-4} < \mathrm{Re}(\bar{c}_{\mu B} - \bar{c}_{\mu W}) < 6.43 \times 10^{-4}\,, \end{gathered} \tag{2.22}$$

$$\begin{gathered} -2.97 \times 10^{-7} < \mathrm{Im}(\bar{c}_{eB} - \bar{c}_{eW}) < 4.51 \times 10^{-7}\,, \\ -0.26 < \mathrm{Im}(\bar{c}_{\mu B} - \bar{c}_{\mu W}) < 0.29\,, \end{gathered} \tag{2.23}$$

where the coefficients are evaluated at the relevant low-energy scale. Notice that the non-vanishing value of $\mathrm{Re}(\bar{c}_{\mu B} - \bar{c}_{\mu W})$ follows from the known $\sim 3.5\sigma$ anomaly in the $(g-2)$ of the muon (see Ref. [27] for an updated review). Among the bounds of Eqs. (2.21), (2.22), (2.23) only those on $\mathrm{Im}(\bar{c}_{eB} - \bar{c}_{eW})$ and $\mathrm{Re}(\bar{c}_{\mu B} - \bar{c}_{\mu W})$ have the sensitivity to probe the values naively expected for these coefficients as reported in Eq. (2.9). In particular, the first one sets an upper bound on $(v/f)^2$ of order 10^{-3} for an $O(1)$ CP phase.

3 Estimates of the effects on physics observables

While the Lagrangian $\Delta\mathcal{L} = \Delta\mathcal{L}_{SILH} + \Delta\mathcal{L}_{F_1} + \Delta\mathcal{L}_{F_2}$ is completely general, the basis of operators of Eqs. (2.2)–(2.4) is particularly useful to characterize the interactions of the

---

[7] The coefficients are evaluated at the following scales: $\mu = m_t$ (Eqs. (2.18) and (2.20)), $\mu = m_W$ (Eq. (2.19)).

Higgs sector. In fact, as already anticipated, one of the main results of Ref. [4] is that of identifying which operators, hence which observables, are sensitive to the strength of the Higgs interactions, rather than merely to the value of the New Physics scale M. In what follows we will discuss this point in greater detail and, starting from the analysis of Refs. [4, 33], we will try to highlight a possible strategy to determine whether the dynamics behind the electroweak breaking is weak or strong. Our analysis will be based on the naive estimates of the Wilson coefficients at the matching scale. In the next Section, we will discuss how the running from the matching scale to the weak scale affects these estimates.

3.1 Operators sensitive to a strongly-interacting Higgs boson

Let us start by considering the effects of the operators O_H, O_T, $O_{u,d,l}$ and O_6: they modify the tree-level couplings of the Higgs boson to fermions, vector bosons and to itself. In the unitary gauge and upon canonical normalization of the Higgs kinetic term, the Lagrangian reads [34]

$$\begin{aligned}\mathcal{L} = &\frac{1}{2}\partial_\mu h\,\partial^\mu h - \frac{1}{2}m_h^2 h^2 - c_3\,\frac{1}{6}\left(\frac{3m_h^2}{v}\right) h^3 + \dots\\ &+ m_W^2\,W_\mu^+ W^{-\,\mu}\left(1 + 2c_W\,\frac{h}{v} + \dots\right) + \frac{1}{2}m_Z^2\,Z_\mu Z^\mu\left(1 + 2c_Z\,\frac{h}{v} + \dots\right)\\ &- \sum_{\psi=u,d,l} m_{\psi^{(i)}}\,\bar{\psi}^{(i)}\psi^{(i)}\left(1 + c_\psi\frac{h}{v} + \dots\right) + \dots\end{aligned} \tag{3.24}$$

where the Higgs couplings $c_{i=W,Z,\psi,3}$, have been defined such that $c_i = 1$ in the SM, and v is defined by Eq. (2.5). Their expressions as functions of the coefficients of the effective Lagrangian (2.2) are given in Table 1. The shifts from the SM value are of order

$$\delta c_i \sim \frac{g_*^2 v^2}{M^2} = \frac{v^2}{f^2}\,. \tag{3.25}$$

Hence, measuring the Higgs couplings probes the strength of its interactions to the new dynamics. Notice that the effective description given by $\Delta\mathcal{L}$ neglects higher powers of (H/f), and is thus valid only if the shifts in the Higgs couplings are small: $\delta c_i \sim (v/f)^2 \ll 1$. If the Higgs doublet is the NG boson of a spontaneously broken symmetry $\mathcal{G} \to \mathcal{H}$, on the other

Higgs couplings	$\Delta\mathcal{L}_{SILH}$	MCHM4	MCHM5
c_W	$1-\bar{c}_H/2$	$\sqrt{1-\xi}$	$\sqrt{1-\xi}$
c_Z	$1-\bar{c}_H/2-2\bar{c}_T$	$\sqrt{1-\xi}$	$\sqrt{1-\xi}$
$c_\psi \;\; (\psi=u,d,l)$	$1-(\bar{c}_H/2+\bar{c}_\psi)$	$\sqrt{1-\xi}$	$\dfrac{1-2\xi}{\sqrt{1-\xi}}$
c_3	$1+\bar{c}_6-3\bar{c}_H/2$	$\sqrt{1-\xi}$	$\dfrac{1-2\xi}{\sqrt{1-\xi}}$
c_{gg}	$8\,(\alpha_s/\alpha_2)\,\bar{c}_g$	0	0
$c_{\gamma\gamma}$	$8\sin^2\theta_W\,\bar{c}_\gamma$	0	0
$c_{Z\gamma}$	$\left(\bar{c}_{HB}-\bar{c}_{HW}-8\,\bar{c}_\gamma\sin^2\theta_W\right)\tan\theta_W$	0	0
c_{WW}	$-2\,\bar{c}_{HW}$	0	0
c_{ZZ}	$-2\left(\bar{c}_{HW}+\bar{c}_{HB}\tan^2\theta_W-4\bar{c}_\gamma\tan^2\theta_W\sin^2\theta_W\right)$	0	0
$c_{W\partial W}$	$-2(\bar{c}_W+\bar{c}_{HW})$	0	0
$c_{Z\partial Z}$	$-2(\bar{c}_W+\bar{c}_{HW})-2\,(\bar{c}_B+\bar{c}_{HB})\tan^2\theta_W$	0	0
$c_{Z\partial\gamma}$	$2\,(\bar{c}_B+\bar{c}_{HB}-\bar{c}_W-\bar{c}_{HW})\tan\theta_W$	0	0

Table 1: The second column reports the values of the Higgs couplings c_i defined in Eq. (3.46) in terms of the coefficients $\bar{c}_i$ of the effective Lagrangian $\Delta\mathcal{L}_{SILH}$. The last two columns show the predictions of the MCHM4 and MCHM5 models in terms of $\xi=(v/f)^2$; the effects of the heavy resonances have been neglected for simplicity, so that only the couplings $c_{W,Z,\psi,3}$ are non-vanishing. The auxiliary parameter α_2 is defined by Eq. (3.43).

hand, it is possible to resum all powers of (H/f) by making use of the invariance under (non-linear) $\mathcal{G}$ transformations. Such an improved effective Lagrangian thus relies only on the expansion in the number of derivatives. For example, in models based on the $SO(5)/SO(4)$ coset [7, 35] the couplings of the Higgs boson to W and Z are predicted to be $c_W = c_Z \equiv c_V = \sqrt{1-\xi}$, where $\xi \equiv (v/f)^2$. The couplings to fermions, on the other hand, are not uniquely fixed by the choice of the coset, but depend on how the SM fermions are coupled to the strong dynamics. The last two columns of Table 1 report the predictions of the Minimal Composite Higgs Model MCHM4 [7] and MCHM5 [35], where the SM fermions couple linearly to composite operators transforming as the spinorial and fundamental representations of $SO(5)$, respectively. For simplicity, the predictions are derived by including only the effects of the Higgs non-linearities, and neglecting those from the heavy resonances, hence only the coefficients c_V, c_ψ and c_3 are non-vanishing. The models MCHM4 and MCHM5 will be considered as benchmarks in the rest of this work.

In general, a shift of the tree-level Higgs couplings of order $(v/f)^2$ implies that the theory gets strongly coupled at energies $\sim 4\pi f$, unless new weakly-coupled physics states set in to regulate the energy growth of the scattering amplitudes. The dominant effect comes from the energy growth of the $V_LV_L \to V_LV_L$ $(V = W^\pm, Z^0)$ scattering amplitudes, which become non-perturbative at the scale $\Lambda_s = 4\pi v/\sqrt{|\bar{c}_H|}$. A modified coupling to the top quark leads instead to strong $V_LV_L \to t\bar{t}$ scattering at energies of order $\Lambda_s = 16\pi^2 v^2/(m_t\sqrt{|\bar{c}_u + \bar{c}_H|})$. The scale of New Physics is thus required to lie below, or at, such ultimate range of validity of the effective theory: $M \lesssim \Lambda_s$.

3.2 Operators sensitive to the scale of New Physics

The operators O_W, O_B can be generated at tree-level by the exchange of heavy particles, for example heavy spin-1 states. In the unitary gauge they are written in terms of the following three operators [8]

$$(D^\mu W^+_{\mu\nu})W^{-\,\nu} h\,, \quad (\partial^\mu Z_{\mu\nu})Z^\nu h\,, \quad (\partial^\mu \gamma_{\mu\nu})Z^\nu h \tag{3.26}$$

---

[8] Here and in the following, derivatives acting on operators in the unitary gauge are covariant under local $U(1)_{em}$ transformations. Operators like $(\partial^\mu Z_{\mu\nu})\gamma^\nu h$ or $(\partial^\mu \gamma_{\mu\nu})\gamma^\nu h$ obviously cannot be generated since they break the $U(1)_{em}$ local symmetry.

plus terms with zero or two Higgs fields. The fact that there are three possible operators in the unitary gauge indicates that their coefficients are related by one identity if the Higgs boson belongs to an $SU(2)$ doublet, see Eq. (3.48). We will discuss this point in greater detail in Section 3.6.

It is easy to see that O_W, O_B give corrections to the tree-level Higgs couplings and generate quartic interactions with one vector boson and two SM fermions that contribute to the three-body decays $h \to VV^* \to V\psi\bar{\psi}$. [9] Indeed, by making use of the equations of motion, [10]

$$iD^\mu W^i_{\mu\nu} = g\, H^\dagger \frac{\sigma^i}{2} \overleftrightarrow{D}_\nu H - ig\, \bar{\psi}\frac{\sigma^i}{2}\gamma_\nu\psi\,, \qquad i\partial^\mu B_{\mu\nu} = \frac{g'}{2}\, H^\dagger \overleftrightarrow{D}_\nu H - ig'\, \bar{\psi}Y\gamma_\nu\psi\,, \tag{3.27}$$

one can rewrite O_W and O_B as

$$O_W = -2\, O_H + \frac{4}{v^2}(H^\dagger H)|D_\mu H|^2 + O'_{Hq} + O'_{HL} \tag{3.28}$$

$$O_B = 2\tan^2\!\theta_W \left(-O_T + O^Y_{H\Psi}\right)\,, \tag{3.29}$$

where the linear combination $O^Y_{H\Psi}$ has been defined in Eq. (2.7). Upon the field redefinition $H \to H - 2\bar{c}_W(H^\dagger H)H/v^2$, the operator $(H^\dagger H)|D_\mu H|^2$ can be rewritten in terms of those in Eq. (2.2). Specifically, Eq. (3.28) becomes: [11]

$$O_W = -6\, O_H + 2\,(O_u + O_d + O_l) - 8\, O_6 + O'_{Hq} + O'_{HL}\,. \tag{3.31}$$

From the estimates of $\bar{c}_W, \bar{c}_B$ and $\bar{c}_H, \bar{c}_T, \bar{c}_\psi, \bar{c}_6$ in Eq. (2.9) one can see that the shifts to the tree-level Higgs couplings due to O_W, O_B are of order $(m_W/M)^2$, hence subdominant in the case of a strongly interacting Higgs boson. Notice that the couplings of the Higgs boson to

---

[9] We thank Riccardo Rattazzi for pointing this out to us.

[10] For simplicity we have left a sum over all fermion representations ψ understood in Eq. (3.27).

[11] By means of Eqs. (3.29) and (3.31) it is thus always possible to remove O_W and O_B provided the coefficients of the other operators are shifted as follows: $\bar{c}_i \to \bar{c}_i + \Delta\bar{c}_i$, with

$$\begin{gathered}\Delta\bar{c}_H = -6\,\bar{c}_W, \quad \Delta\bar{c}_T = -2\tan^2\!\theta_W\,\bar{c}_B\,, \quad \Delta\bar{c}_6 = -8\,\bar{c}_W, \quad \Delta\bar{c}_\psi = 2\,\bar{c}_W \\ \Delta\bar{c}'_{Hq} = \Delta\bar{c}'_{HL} = \bar{c}_W \\ 6\,\Delta\bar{c}_{Hq} = \frac{3}{2}\,\Delta\bar{c}_{Hu} = -3\,\Delta\bar{c}_{Hd} = -2\,\Delta\bar{c}_{HL} = -\Delta\bar{c}_{Hl} = -2\tan^2\!\theta_W\,\bar{c}_B\,.\end{gathered} \tag{3.30}$$

W and Z get different shifts from O_B (since $\Delta\bar{c}_T \neq 0$). In practice, the constraint (2.12) bounds this custodial-symmetry breaking effect down to an unobservable level, unless some fine tuning is in place in the combination $\bar{c}_W + \bar{c}_B$ so that $\bar{c}_B$ can be large. Notice that despite the operator O_T is generated after using the equations of motion, its contribution to $\Delta\epsilon_1$ (corresponding to a non-vanishing $\hat{T}$ parameter [14, 12]) is exactly canceled by the vertex correction implied by the linear combination of fermionic operators which is also generated. [12] This is of course expected, since O_W, O_B only contribute to ϵ_3, and not to ϵ_1.

In general, the contribution of O_W, O_B to inclusive observables, in particular to the Higgs decay rates, is of order (m_W^2/M^2):

$$\frac{\delta\Gamma(h \to VV)}{\Gamma(h \to VV)}\bigg|_{O_W,O_B} \sim O\!\left(\frac{m_W^2}{M^2}\right) , \tag{3.32}$$

where in this case $VV = W^{(*)}W^*, Z^{(*)}Z^*, Z^{(*)}\gamma, \gamma\gamma$. This implies that these operators are sensitive only to the value of the scale of New Physics M, and do not probe the coupling strength g_*. From the quantitative side, the constraint (2.12) suggests that their effects in inclusive Higgs decay rates is too small to be observable. For example, we find that for small $\bar{c}_{W,B}$ the tree-level correction to the WW and ZZ partial rates is well approximated by: [13]

$$\frac{\Gamma(h \to W^{(*)}W^*)}{\Gamma(h \to W^{(*)}W^*)_{SM}} \simeq 1 + 2.2\,\bar{c}_W , \qquad \frac{\Gamma(h \to Z^{(*)}Z^*)}{\Gamma(h \to Z^{(*)}Z^*)_{SM}} \simeq 1 + 2.0\left(\bar{c}_W + \tan^2\!\theta_W\,\bar{c}_B\right) . \tag{3.33}$$

Notice that despite its custodial invariance, the operator O_W affects in a slightly different way the decay of the Higgs boson into WW and ZZ, due to the fact that at least one of the two final vector bosons is off-shell. [14] At the one-loop level O_W also contributes to the Higgs

---

[12] See for example Eq. (9.10) of Ref. [13].

[13] Here and in the following our approximated formulas have been obtained by setting $m_h = 125\,\text{GeV}$ and fully including the QCD corrections. The EW corrections to the decay rates are instead not included, since their effect on the numerical prefactor appearing in front of the coefficients $\bar{c}_i$ is of order $(v^2/f^2)(\alpha_2/4\pi)$ and thus beyond the accuracy of our computation. See Appendix E.6 for more details.

[14] It is easy to check that for $m_h > 2m_Z$ and on-shell decays one has:

$$\frac{\Gamma(h \to WW)}{\Gamma(h \to WW)_{SM}} \simeq 1 + 4\,\bar{c}_W , \qquad \frac{\Gamma(h \to ZZ)}{\Gamma(h \to ZZ)_{SM}} \simeq 1 + 4\left(\bar{c}_W + \tan^2\!\theta_W\,\bar{c}_B\right) . \tag{3.34}$$

These formulas coincide with those of Eqs. (79)–(80) of Ref. [4], which are thus valid only for on-shell decays.

decays into $Z\gamma$ and $\gamma\gamma$ (while O_B does not). We find:

$$\frac{\Gamma(h \to Z\gamma)}{\Gamma(h \to Z\gamma)_{SM}} \simeq 1 + 4.2\,\bar{c}_W\,, \qquad \frac{\Gamma(h \to \gamma\gamma)}{\Gamma(h \to \gamma\gamma)_{SM}} \simeq 1 + 5.0\,\bar{c}_W\,, \tag{3.35}$$

which agree with Eqs. (82) and (83) of Ref. [4]. [15] For $\bar{c}_{W,B} \sim 10^{-3}$ the above approximate formulas imply corrections too small to be observed at the LHC. On the other hand, one could try to take advantage of the different predictions in terms of angular and invariant mass distributions which are implied by the dimension-6 operators compared to the tree-level SM prediction. The most promising strategy could be in fact that based on the analysis of the angular distributions of the final fermions [36–38]. In the ideal case in which one is able to kill completely the SM tree-level contribution by means of appropriate kinematic cuts, the relative effect of NP becomes of order

$$\frac{d\Gamma(h \to VV)}{d\Omega} \Big/ \left(\frac{d\Gamma(h \to VV)}{d\Omega}\right)_{SM} \lesssim 1 + \bar{c}_{W,B}\,\frac{16\pi^2}{g^2}\,, \tag{3.36}$$

which might leave room for observable effects even for $\bar{c}_{W,B} \sim O(10^{-3})$. Clearly, a more precise assessment of the efficiency of such a strategy requires a dedicated analysis [39].

3.3 Operators generated at the one-loop level

Let us now focus on the operators O_{HW}, O_{HB}, O_γ and O_g, which are generated at the one-loop level. In the unitary gauge, $O_{HW,HB,\gamma}$ are rewritten in terms of

$$W^+_{\mu\nu}W^{-\,\mu\nu}h\,, \quad Z_{\mu\nu}Z^{\mu\nu}h\,, \quad \gamma_{\mu\nu}\gamma^{\mu\nu}h\,, \quad Z_{\mu\nu}\gamma^{\mu\nu}h \tag{3.37}$$

plus other terms with zero or two Higgs fields. Since the coefficients of the above four operators are functions of $\bar{c}_{HW}$, $\bar{c}_{HB}$ and $\bar{c}_\gamma$, they are related by one identity, see Eq. (3.47). We will discuss this point in greater detail in Section 3.6.

As implied from the naive estimates (2.9), the contribution of $O_{HW,HB,\gamma}$ to the WW and ZZ inclusive rates is of order ($VV = WW, ZZ$)

$$\left.\frac{\delta\Gamma(h \to VV)}{\Gamma(h \to VV)}\right|_{O_\gamma, O_{HW}, O_{HB}} \sim O\!\left(\frac{m_W^2}{16\pi^2 f^2}\right)\,. \tag{3.38}$$

---

[15] The easiest way to compute the one-loop contribution of O_W to the $Z\gamma$ and $\gamma\gamma$ rates is by using Eq. (3.28) to rewrite this operator in terms of the others. Among the operators generated in this way, only O_H gives a contribution. Notice that if Eq. (3.31) is used instead, one has to take into account also the contribution of $(O_u + O_d + O_l)$ and the shift to the Fermi constant induced by $O'_{Hq} + O'_{HL}$.

Although such an effect depends on the Higgs interaction strength, it is suppressed compared to Eq. (3.32) by a loop factor. We find that the following approximate formulas hold [16]

$$\begin{aligned}\frac{\Gamma(h\to W^{(*)}W^{*})}{\Gamma(h\to W^{(*)}W^{*})_{SM}} &\simeq 1+3.7\,\bar c_{HW}\,,\\ \frac{\Gamma(h\to Z^{(*)}Z^{*})}{\Gamma(h\to Z^{(*)}Z^{*})_{SM}} &\simeq 1+3.0\left(\bar c_{HW}+\tan^2\!\theta_W\,\bar c_{HB}\right)-0.26\,\bar c_\gamma\,.\end{aligned}\tag{3.40}$$

While the contribution due to $\bar c_{HB}$ and $\bar c_\gamma$ explicitly violates the custodial symmetry and thus differentiates WW from ZZ, the different numerical factor multiplying $\bar c_{HW}$ in the two formulas above is due to the off-shellness of at least one of the two vector bosons, similarly to Eq. (3.33). Although there is currently no stringent bound on the coefficients $\bar c_{HW,HB,\gamma}$, the estimate (2.9) suggests that their correction to inclusive rates is unobservable at the LHC. As discussed in the previous section, on the other hand, a study of the angular and invariant mass distributions of these decays can potentially uncover the effect of New Physics. In particular, an estimate similar to that of Eq. (3.36) can be derived also for $\bar c_{HW,HB,\gamma}$.

The processes $h\to\gamma\gamma$, $h\to Z\gamma$ and $h\to gg$ (or equivalently $gg\to h$) can in principle test the Higgs interaction strengths much more powerfully, since they arise at the one-loop level in the SM. Naively one expects:

$$\left.\frac{\delta\Gamma(h\to gg,\gamma\gamma,Z\gamma)}{\Gamma(h\to gg,\gamma\gamma,Z\gamma)}\right|_{O_g,O_\gamma,O_{HW},O_{HB}} \sim O\!\left(\frac{v^2}{f^2}\right)\,. \tag{3.41}$$

---

[16] For $m_h > 2m_Z$ and on-shell decays, we find instead

$$\begin{aligned}\frac{\Gamma(h\to WW)}{\Gamma(h\to WW)_{SM}} &\simeq 1+8\,\bar c_{HW}\,,\\ \frac{\Gamma(h\to ZZ)}{\Gamma(h\to ZZ)_{SM}} &\simeq 1+8\,(\bar c_{HW}+\tan^2\theta_W\,\bar c_{HB})-16\tan^2\theta_W\sin^2\theta_W\,\bar c_\gamma\,.\end{aligned}\tag{3.39}$$

Comparing with the analog formulas in Eqs. (79) and (80) of Ref. [4], we find that in these latter there is a missing factor 2 and the term proportional to $\bar c_\gamma$ was not included either. Notice also that the effect of the off-shellness of the gauge bosons is rather large, as one can see by comparing Eq. (3.39) with Eq. (3.40).

We find that the following approximate formulas hold to good accuracy for small $\bar{c}_i$'s:

$$\begin{aligned}
\frac{\Gamma(h \to gg)}{\Gamma(h \to gg)_{SM}} &\simeq 1 + 22.2\, \bar{c}_g \, \frac{4\pi}{\alpha_2} \\
\frac{\Gamma(h \to \gamma\gamma)}{\Gamma(h \to \gamma\gamma)_{SM}} &\simeq 1 - 0.54\, \bar{c}_\gamma \, \frac{4\pi}{\alpha_{em}} \,, \\
\frac{\Gamma(h \to Z\gamma)}{\Gamma(h \to Z\gamma)_{SM}} &\simeq 1 + 0.19 \left(\bar{c}_{HW} - \bar{c}_{HB} + 8\, \bar{c}_\gamma \sin^2\theta_W \right) \frac{4\pi}{\sqrt{\alpha_2 \alpha_{em}}} \,,
\end{aligned} \tag{3.42}$$

where we have conveniently defined

$$\alpha_2 \equiv \frac{\sqrt{2}\, G_F m_W^2}{\pi} \,, \tag{3.43}$$

and by α_{em} we indicate the value of the running electromagnetic coupling $\alpha_{em}(q^2 = 0)$ in the Thomson limit. If the Higgs boson is a NG boson, the coefficients $\bar{c}_g$ and $\bar{c}_\gamma$ are further suppressed by a factor $(g_{\not{G}}/g_*)^2$, see Eq. (2.10), where $g_{\not{G}}$ is a weak coupling. This implies that in this class of theories the corrections to $\Gamma(h \to \gamma\gamma)$ and $\Gamma(h \to gg)$ depend only on the scale of New Physics and not on the Higgs interaction strength. In fact, in the case of minimal models with linear couplings, like for example the MCHM4 and MCHM5, the low energy theorem [40, 41] implies that the leading contribution to the $\gamma\gamma$ and gg decay rates from the virtual exchange of heavy fermions is additionally suppressed [42–45] due to a cancellation between the effect parametrized by $\bar{c}_{g,\gamma}$ and the one that follows from the shift in the top Yukawa coupling due to $\bar{c}_u$ and $\bar{c}_H$ (see Ref. [44] for an interesting exception). In general, in theories with a pNGB Higgs boson the local corrections to the rates $\Gamma(h \to \gamma\gamma)$ and $\Gamma(h \to gg)$ from O_γ and O_g are expected to be small and subdominant compared to the effect from the modified tree-level Higgs couplings.

3.4 Fermionic operators

The fermionic operators in $\Delta\mathcal{L}_{F_1}$ are sensitive to the strength of the couplings of the Higgs boson and of the SM fermions to the new dynamics. They lead to contact corrections to the three-body decays $h \to VV^* \to V\psi\psi$ which are naively of order

$$\frac{\delta\Gamma(h \to V\bar{\psi}\psi)}{\Gamma(h \to V\bar{\psi}\psi)} \sim O\!\left(\frac{v^2}{f^2} \frac{\lambda_\psi^2}{g_*^2} \right) . \tag{3.44}$$

Compared to the corrections from O_W and O_B, the effect of the fermionic operators is potentially enhanced by a factor (λ_ψ^2/g^2). In practice, the possibility of large fermionic couplings λ_ψ is strongly constrained by LEP, see Eqs. (2.14)-(2.16). The only scenarios in which a large degree of compositeness of either the left- or right-handed quarks is not ruled out are those in which the corresponding operators in $\Delta\mathcal{L}_{F_1}$ are not generated as due to some protecting symmetry (see for example Refs. [21, 46, 47]). Large corrections to the inclusive rate of the three-body decays $h \to V\bar{\psi}\psi$ from $\Delta\mathcal{L}_{F_1}$ are thus excluded, while the possibility of detecting the effects of these operators through the analysis of differential distributions should be explored, similarly to what has been discussed for O_W and O_B.

Among the dipole operators in $\Delta\mathcal{L}_{F_2}$, those with light fermions are already strongly constrained by current precision data, but potentially sizable effects could still come from the operators involving the top quark. For example, the contribution of O_{tG} to $gg \to h$, $gg \to t\bar{t}$, $gg \to t\bar{t}h$ is of order $E^2/(16\pi^2 f^2)$, where E is the energy scale relevant in the process. More in detail

$$\frac{\delta\sigma(gg \to h)}{\sigma(gg \to h)} \sim \hat{c}_{tG}\,, \qquad \frac{\delta\sigma(gg \to t\bar{t})}{\sigma(gg \to t\bar{t})} \sim \hat{c}_{tG}\,\frac{\sqrt{s}}{m_t}\,, \qquad \frac{\delta\sigma(gg \to t\bar{t}h)}{\sigma(gg \to t\bar{t}h)} \sim \hat{c}_{tG}\,\frac{s}{m_t^2}\,, \tag{3.45}$$

where we have defined $\hat{c}_{tG} \equiv \mathrm{Re}(\bar{c}_{tG})\,(m_t^2/m_W^2) \sim m_t^2/(16\pi^2 f^2) \simeq 3 \times 10^{-3}(v^2/f^2)$. Notice that the experimental limit on the neutron EDM puts an upper bound on the imaginary part of $\hat{c}_{tG}$ at the 10^{-4} level, see Eq. (2.18), which indicates that this is currently the most sensitive experiment on $\mathrm{Im}(\bar{c}_{tG})$. Some mechanism is however required to suppress the imaginary parts of the dipole operators involving light fermions, in order to satisfy the stringent constraints of Eq. (2.17). By the same mechanism also $\mathrm{Im}(\bar{c}_{tG})$ could be suppressed, so that the processes of Eq. (3.45) are essential to probe the contribution of O_{tG} due to $\mathrm{Re}(\bar{c}_{tG})$. From Eq. (3.45) and the naive estimate of $\hat{c}_{tG}$ it follows that the most sensitive process is perhaps $gg \to t\bar{t}$, in particular the events at large invariant mass, although a precision larger than the one currently achieved is required to constrain (v/f). To this aim, the analysis of differential distributions and spin correlations could be a successful strategy [48, 25]. The NP contribution to the process $gg \to t\bar{t}h$ can in principle get the largest enhancement from a cut on $\sqrt{s}$, but the small rate might limit the actual sensitivity achievable at the LHC [49]. Finally, additional information comes from the experimental limits on top anomalous couplings obtained at the Tevatron and the LHC, although their sensitivity on NP is expected to be much smaller by naive estimate. The operator O_{tW}, in particular, gives the largest effect and generates the anomalous coupling $g_R(g/m_W)\bar{b}_L\sigma^{\mu\nu}W^-_{\mu\nu}t_R$ [26]. Naively one expects $g_R = (4m_t/m_W)\,\bar{c}_{tW} \sim m_t m_W/(16\pi^2 f^2) = 1.5 \times 10^{-3}\,(v/f)^2$, an effect too small to be observed even for f of order v.

3.5 Non-linear Lagrangian for a Higgs-like scalar

Summarizing, by working in the unitary gauge and in the basis of fermion mass eigenstates, the effective Lagrangian relevant for Higgs physics reads as follows [34]

$$
\begin{aligned}
\mathcal{L} = {} & \frac{1}{2}\partial_\mu h\; \partial^\mu h - \frac{1}{2}m_h^2 h^2 - c_3\,\frac{1}{6}\left(\frac{3m_h^2}{v}\right) h^3 - \sum_{\psi=u,d,l} m_{\psi^{(i)}}\;\bar{\psi}^{(i)}\psi^{(i)}\left(1 + c_\psi \frac{h}{v} + \dots\right) \\
& + m_W^2\, W_\mu^+ W^{-\,\mu}\left(1 + 2c_W\,\frac{h}{v} + \dots\right) + \frac{1}{2}m_Z^2\, Z_\mu Z^\mu \left(1 + 2c_Z\,\frac{h}{v} + \dots\right) + \dots \\
& + \left(c_{WW}\, W^+_{\mu\nu}W^{-\mu\nu} + \frac{c_{ZZ}}{2}\, Z_{\mu\nu}Z^{\mu\nu} + c_{Z\gamma}\, Z_{\mu\nu}\gamma^{\mu\nu} + \frac{c_{\gamma\gamma}}{2}\,\gamma_{\mu\nu}\gamma^{\mu\nu} + \frac{c_{gg}}{2}\, G^a_{\mu\nu}G^{a\mu\nu}\right)\frac{h}{v} \\
& + \left(c_{W\partial W}\left(W^-_\nu D_\mu W^{+\mu\nu} + h.c.\right) + c_{Z\partial Z}\, Z_\nu\partial_\mu Z^{\mu\nu} + c_{Z\partial\gamma}\, Z_\nu\partial_\mu\gamma^{\mu\nu}\right)\frac{h}{v} + \dots
\end{aligned}
\tag{3.46}
$$

where, we recall, v is defined in Eq. (2.5). We have shown only terms involving up to three bosonic fields, and we have omitted in particular those involving fermions that follow from $\Delta\mathcal{L}_{F_1} + \Delta\mathcal{L}_{F_2}$. Their form can be easily derived from Eqs. (2.3) and (2.4). The relations between the couplings appearing in Eq. (3.46) and the coefficients of the dimension-6 operators in Eq. (2.2) are reported in Table 1. It is worth noting that the same Lagrangian (3.46) applies also to the case in which the electroweak symmetry $SU(2)_L \times U(1)_Y$ is non-linearly realized and h is a generic CP-even scalar, singlet of the custodial symmetry, not necessarily connected with the EW symmetry breaking. Indeed, each of the terms in (3.46), being invariant under local $U(1)_{em}$ transformations, can be dressed up with the Nambu-Goldstone bosons that are eaten to form the longitudinal W and Z polarizations and made manifestly $SU(2)_L \times U(1)_Y$ gauge invariant [50] (see also Ref. [51]). The explicit expression in such a basis has been given in Refs. [52,53] at the level of four-derivative operators. In this sense the effective Lagrangian (3.46) is a generic tool to understand the origin of the newly discovered boson and the role it plays in the electroweak symmetry breaking dynamics. It is valid for arbitrary values of the couplings c_i appearing in Eq. (3.46), and it can be used to make computations of observable quantities at a given order in an expansion in E/M and in $\alpha_{SM}/4\pi$, where by the latter we indicate the generic SM loop expansion parameter. That is in full analogy with other well-known effective theories, see Ref. [3]. It should be stressed that, according to a well established methodology and similarly to Eq. (2.2), in this effective Lagrangian all quantum fluctuations associated to short-length modes (high-energy modes) have already been considered and are parametrized by local operators with an increasing number of derivatives, while quantum fluctuations (loop diagrams) involving the light modes still have to be taken into account. For instance, top loops will give an additional contribution to the on-shell h-gluon-gluon coupling. While Eq. (3.46) is general, the effective Lagrangian (2.2) assumes that h is part of an $SU(2)_L$ doublet and further relies on the expansion in powers of H/f. As such, it is valid only in the limit of small deviations of the Higgs couplings from their SM values and up to corrections of order $O(v^2/f^2)$.

3.6 Implications of custodial symmetry

Another difference between the non-linear Lagrangian (3.46) and the SILH Lagrangian (2.2) is that the first one contains two more free parameters. This means that there are two relations among the couplings of Eq. (3.46) which hold at the level of dimension-6 operators if the Higgs is part of a doublet. As noticed in Sections (3.2) and (3.3), the first identity relates c_{WW}, c_{ZZ}, $c_{Z\gamma}$ and $c_{\gamma\gamma}$, while the second relates $c_{W\partial W}$, $c_{Z\partial Z}$ and $c_{Z\partial\gamma}$. They read:

$$c_{WW} - c_{ZZ}\cos^2\theta_W = c_{Z\gamma}\sin 2\theta_W + c_{\gamma\gamma}\sin^2\theta_W \tag{3.47}$$

$$c_{W\partial W} - c_{Z\partial Z}\cos^2\theta_W = \frac{c_{Z\partial\gamma}}{2}\sin 2\theta_W\,. \tag{3.48}$$

In fact both identities are not special to the case in which the Higgs is a doublet, but are a general consequence of custodial symmetry. This latter is accidental in the SILH Lagrangian if one restricts to the operators that lead to derivative couplings of the Higgs to vector bosons. Starting at the dimension-8 order, it is possible to write cutodial-breaking operators that lead to couplings that violate the relations (3.47) and (3.48). For instance

$$\frac{\bar{c}_{8WW}\, g^2}{m_W^2 v^2}\left(H^\dagger W^a_{\mu\nu}\sigma^a H\right)\left(H^\dagger W^{b\,\mu\nu}\sigma^b H\right) + \frac{i\bar{c}_{8W}\, g}{v^2 m_W^2}\left(H^\dagger\sigma^a H\right)\left(D^\mu W_{\mu\nu}\right)^a\left(H^\dagger \overleftrightarrow{D^\nu} H\right) \tag{3.49}$$

gives rise to

$$\begin{gathered} c_{Z\partial Z} = -4\bar{c}_{8W}, \qquad c_{Z\partial\gamma} = -4\tan\theta_W \bar{c}_{8W} \,, \\ c_{ZZ} = 8\cos^2\theta_W \bar{c}_{8WW} \,, \qquad c_{Z\gamma} = 4\sin 2\theta_W \bar{c}_{8WW} \,, \qquad c_{\gamma\gamma} = 8\sin^2\theta_W \bar{c}_{8WW} \,, \end{gathered} \tag{3.50}$$

and the relations (3.47) and (3.48) are not fulfilled. [17]

A third relation holds on the non-derivative couplings c_W and c_Z if one assumes that custodial symmetry is an invariance of the Lagrangian (2.2), so that $\bar{c}_T = 0$; it reads:

$$c_W = c_Z \,. \tag{3.51}$$

As said above, while all three identities (3.47), (3.48) and (3.51) are a consequence of custodial symmetry, the first two are accidental at the level of dimension-6 operators if the Higgs is part of a doublet.

To show that Eqs. (3.47), (3.48) and (3.51) follow from custodial invariance, let us consider the case in which the EWSB dynamics has a global $SU(2)_L \times SU(2)_R$ symmetry, and imagine to fully gauge this group by enlarging the hypercharge to a whole triplet of $SU(2)_R$. In this case the diagonal custodial $SU(2)_V$ is exact even though $g' \neq g$. The left and right gauge fields couple to the conserved currents of $SU(2)_L \times SU(2)_R$ and the interactions among two gauge fields and the Higgs boson are fully characterized in momentum space by three form factors:

$$(\Gamma_{LL})^{\mu\nu}_{ij}(p_1,p_2)\, L^i_\mu L^j_\nu h + (\Gamma_{LR})^{\mu\nu}_{ij}(p_1,p_2)\, L^i_\mu R^j_\nu h + (\Gamma_{RR})^{\mu\nu}_{ij}(p_1,p_2)\, R^i_\mu R^j_\nu h \,. \tag{3.52}$$

Here p_1, p_2 are the momenta of the gauge fields and each form factor can be computed in terms of a Green function with two conserved currents, $\Gamma^{\mu\nu}_{ik} = \langle J^\mu_i J^\nu_k | h\rangle$. In addition to the usual massive W and Z bosons, which form a triplet $\hat{V}^i_\mu$ of the custodial group, in this case there is a whole triplet of massless $SU(2)_V$ gauge fields (the photon plus its charged companion), V^i_μ. The mass eigenstates V_μ and $\hat{V}_\mu$ are related to the left and right gauge fields through a rotation by an angle θ_W, where $\tan\theta_W = g'/g$. Their cubic interactions with the Higgs boson are thus characterized by three form factors, which are linear combinations of those in Eq. (3.52):

---

[17] The two operators in (3.49) give rise to the oblique parameter $\hat{U}$, see for instance Ref. [12]: $\hat{U} = -\bar{c}_{8W} - 2\bar{c}_{8HW}$ while $\hat{S} = \bar{c}_{8HW}$.

$$\begin{aligned}
\Gamma_{VV} &= \sin^2\!\theta_W\,\Gamma_{LL} + \frac{\sin 2\theta_W}{2}\,(\Gamma_{LR}+\Gamma_{RL}) + \cos^2\!\theta_W\,\Gamma_{RR} \\
\Gamma_{\hat{V}V} &= \frac{\sin 2\theta_W}{2}\Gamma_{LL} + \big(\cos^2\!\theta\,\Gamma_{LR} - \sin^2\!\theta\,\Gamma_{RL}\big) - \frac{\sin 2\theta_W}{2}\Gamma_{RR} \\
\Gamma_{\hat{V}\hat{V}} &= \cos^2\!\theta_W\,\Gamma_{LL} - \frac{\sin 2\theta_W}{2}\,(\Gamma_{LR}+\Gamma_{RL}) + \sin^2\!\theta_W\,\Gamma_{RR}\,,
\end{aligned} \tag{3.53}$$

where we have defined $\Gamma^{\mu\nu}_{RL}(p_1,p_2) \equiv \Gamma^{\nu\mu}_{LR}(p_2,p_1)$. Notice, in particular, that in this case the same form factor $\Gamma_{\hat{V}\hat{V}}$ describes the interaction of two W's and two Z's to the Higgs boson, as due to custodial invariance.

The physical limit where only $SU(2)_L \times U(1)_Y$ is gauged is obtained by simply switching off the unphysical $R^{1,2}_\mu$ fields. The interactions of two neutral vector bosons to the Higgs are still described by the relations of Eq. (3.53), where $\Gamma_{ZZ} = \Gamma_{\hat{V}\hat{V}}$, $\Gamma_{\gamma\gamma} = \Gamma_{VV}$ and $\Gamma_{Z\gamma} = \Gamma_{\hat{V}V}$. In the charged sector, instead, the W corresponds to a pure left gauge field, since it has no mixing with right-handed ones. This implies that its form factor is given by the last formula of Eq. (3.53) with $\theta_W = 0$, that is: $\Gamma_{WW} = \Gamma_{LL}$. The four physical form factors are linear combinations of the three defined in Eq. (3.52), and are thus related by one identity:

$$\Gamma^{\mu\nu}_{WW}(p_1,p_2) - \Gamma^{\mu\nu}_{ZZ}(p_1,p_2)\cos^2\!\theta_W = \big(\Gamma^{\mu\nu}_{Z\gamma}(p_1,p_2) + \Gamma^{\nu\mu}_{Z\gamma}(p_2,p_1)\big)\,\frac{\sin 2\theta_W}{2} + \Gamma^{\mu\nu}_{\gamma\gamma}(p_1,p_2)\sin^2\!\theta_W\,. \tag{3.54}$$

Notice that this relation is a consequence of our initial assumption of $SU(2)_L \times SU(2)_R$ invariance of the EWSB dynamics. The custodial $SU(2)_V$ is broken in this case only by the gauging of hypercharge. For $g' = 0$ the custodial symmetry is unbroken and Eq. (3.54) implies $\Gamma_{WW} = \Gamma_{ZZ}$. It is straightforward to derive the relations (3.47), (3.48) and (3.51) from Eq. (3.54). At quadratic order in the momenta, the form factors can be computed from the effective Lagrangian (3.46); one has:

$$\begin{aligned}
\Gamma^{\mu\nu}_{WW}(p_1,p_2) &= 2m_W^2 c_W\,\eta^{\mu\nu} - 2c_{WW}P^{\mu\nu}_{12} - c_{W\partial W}\,(P^{\mu\nu}_1 + P^{\mu\nu}_2) \\
\Gamma^{\mu\nu}_{ZZ}(p_1,p_2) &= 2m_Z^2 c_Z\,\eta^{\mu\nu} - 2c_{ZZ}P^{\mu\nu}_{12} - c_{Z\partial Z}\,(P^{\mu\nu}_1 + P^{\mu\nu}_2) \\
\Gamma^{\mu\nu}_{Z\gamma}(p_1,p_2) &= -\,2c_{Z\gamma}P^{\mu\nu}_{12} - c_{Z\partial\gamma}P^{\mu\nu}_2 \\
\Gamma^{\mu\nu}_{\gamma\gamma}(p_1,p_2) &= -\,2c_{\gamma\gamma}P^{\mu\nu}_{12}\,,
\end{aligned} \tag{3.55}$$

where we have defined $P^{\mu\nu}_1 \equiv \eta^{\mu\nu}p_1^2 - p_1^\mu p_1^\nu$, $P^{\mu\nu}_2 \equiv \eta^{\mu\nu}p_2^2 - p_2^\mu p_2^\nu$ and $P^{\mu\nu}_{12} \equiv \eta^{\mu\nu}p_1{\cdot}p_2 - p_1^\nu p_2^\mu$. This is in fact the most general decomposition which follows at the $O(p^2)$ level for an on-shell

Higgs boson by assuming CP invariance and requiring that: *i)* the Γ_{WW}, Γ_{ZZ} and $\Gamma_{\gamma\gamma}$ form factors are symmetric under the exchange $\{p_1, \mu\} \leftrightarrow \{p_2, \nu\}$; *ii)* the $\Gamma_{\gamma\gamma}$ and $\Gamma_{Z\gamma}$ form factors satisfy the Ward identities implied by $U(1)_{em}$ local invariance:

$$p_{1\mu}\Gamma^{\mu\nu}_{\gamma\gamma}(p_1, p_2) = 0 = p_{2\nu}\Gamma^{\mu\nu}_{\gamma\gamma}(p_1, p_2)\,, \qquad p_{2\nu}\Gamma^{\mu\nu}_{Z\gamma}(p_1, p_2) = 0\,. \tag{3.56}$$

Additional structures proportional to $p_{1\mu}$ and $p_{2\nu}$ can be omitted since they give vanishing contributions both when the vector bosons are on-shell and when they decay into a pair of fermions by coupling to the corresponding conserved current. Inserting Eq. (3.55) into (3.54) one then obtains the identities (3.47), (3.48) and (3.51).

From the above discussion it follows that if custodial symmetry is an invariance of the EWSB dynamics, the effective Lagrangians (3.46) and (2.2) have the same number of free parameters, in terms of which all observables can be computed. This is true also if one considers the fermionic operators (for a Higgs doublet these are listed in Eqs. (2.3) and (2.4)), as long as one focuses on terms with one Higgs boson. This means that by using single-Higgs processes alone, one cannot distinguish the case in which the Higgs boson is part of a doublet from the more general situation. The only possible strategy to this aim is exploiting the connection among processes with zero, one and two Higgs bosons which is implied by the Lagrangian (2.1) at $O(v^2/f^2)$ and does not hold in the case of the more general non-linear Lagrangian. As a consequence of such connection, the bounds that EW and flavor data set on operators with zero Higgs fields severely constrain the size of the NP effects in Higgs processes, as discussed in Section (2.1). If one were to find that single-Higgs processes violate these constraints, this would be an indication that the Higgs is not part of a doublet. Furthermore, processes with double Higgs boson production can be predicted to a certain extent in terms of single-Higgs couplings, and can thus be used to probe the nature of the Higgs boson [54].

4 Implementing the Higgs effective Lagrangian beyond the tree level

In this section we address a few issues related to the use of the effective Lagrangians (2.1) and (3.46) beyond the tree level, as required to make Higgs precision physics without assuming the validity of the Standard Model. While the methodology is well established and various examples of its application exist in several different contexts, we think that a dedicated discussion can be useful to better clarify some specific points (see also Ref. [55] for a recent discussion). As an illustrative though important example, we will consider the calculation of the Higgs partial decay widths, and show how the corrections from dimension-6 operators can be incorporated in a consistent way. As a by-product of our analysis and to better demonstrate its applicability, we present in Appendix E a modified version of the program `HDECAY` [56] which features a full implementation of the effective Lagrangian $\Delta\mathcal{L}_{SILH}$, Eq. (2.2), as well as its generalization to the case of a non-linearly realized EW symmetry, Eq. (3.46).

A first difficulty which arises when using either Eq. (2.1) or (3.46) is the presence of multiple expansion parameters. For generic values of the Higgs couplings c_i, the validity of the effective Lagrangian (3.46) is based on a double perturbative expansion in the SM couplings, $\alpha_{SM}/4\pi$, and in powers of E/M. The effective Lagrangian (2.1) further assumes $(v/f) \ll 1$, which implies small shifts in the Higgs couplings: $c_i = 1+\delta c_i$, with $\delta c_i \lesssim O(v^2/f^2)$. All these expansion parameters must be properly taken into account when performing calculations. Furthermore, the non-renormalizability of the effective theory implies the presence of additional divergences compared to the SM case which must be absorbed by a renormalization of the Wilson coefficients of local operators.

4.1 RG evolution of the Wilson coefficients

Let us discuss the issue of the renormalization and RG evolution of the Wilson coefficients first. As done in the previous sections, we will assume that the Higgs boson is part of an $SU(2)_L$ doublet and use the Lagrangian (2.1). Since we are only interested in the divergent structure of the diagrams, it is convenient to work in the limit of unbroken $SU(2)_L \times U(1)_Y$ and compute the Green functions in terms of the Higgs doublet H. The only 1-loop diagrams which generate additional logarithmic divergences are those featuring one insertion of the effective vertices from dimension-6 operators. By dimensional analysis, further insertions of the effective vertices lead to power-divergent contributions to dimension-6 operators (which are irrelevant to determine the RG running) and log-divergent contributions to higher-dimensional operators. The same counting holds also at higher loop level: the only log-divergent contribution to dimension-6 operators comes from diagrams with one insertion of the effective couplings, and is thus suppressed by extra powers of the SM expansion parameter $\alpha_{SM}/4\pi$. This is in analogy with the renormalization of the pion effective Lagrangian in the chiral limit, see Ref. [57]. It thus follows that the RG equation is linear and homogeneous in the $\bar{c}_i$, and different operators with the same quantum numbers will in general mix with each other. At leading order in α_{SM}, with $\alpha_{SM} = \alpha_{em}, \alpha_2, \alpha_s$, respectively, in the case of electromagnetic, weak and QCD corrections, one has

$$\bar{c}_i(\mu) = \left(\delta_{ij} + \gamma_{ij}^{(0)} \frac{\alpha_{SM}(\mu)}{4\pi} \log\Big(\frac{\mu}{M}\Big)\right) \bar{c}_j(M)\,, \tag{4.57}$$

where $\gamma_{ij}^{(0)}$ is the leading-order coefficient of the anomalous dimension. Some elements of the anomalous dimension matrix $\gamma_{ij}^{(0)}$ have been recently computed in Refs. [58, 59].

In the case in which the Higgs boson and possibly the SM quarks (in particular the top and the bottom) are strongly coupled to the new dynamics, the leading RG running effect comes from loops of these particles and can be as large as $\Delta\bar{c}_i/\bar{c}_i(M) \sim (g_*^2/16\pi^2)\log(M/\mu)$ or $(\lambda_\psi^2/16\pi^2)\log(M/\mu)$. This must be compared to the effects of order $(g_{SM}^2/16\pi^2)\log(M/\mu)$ from loops of gauge fields. For example, the insertion of $\bar{c}_H$ in the diagram (a) of Fig. 1 leads to a renormalization of $O_{W+B} \equiv O_W + O_B$:

$$\bar{c}_{W+B}(\mu) = \bar{c}_{W+B}(M) - \frac{1}{6}\,\frac{\alpha_2}{4\pi}\log\left(\frac{\mu}{M}\right)\bar{c}_H(M)\,, \tag{4.58}$$

where α_2 has been defined in Eq. (3.43). It is well known that this RG running is associated with the IR contribution to the ϵ_3 parameter, and the same coefficient $\gamma_{W+B,H}^{(0)} = -1/6$ can indeed be extracted from self-energy diagrams [60]. From the estimates of Eq. (2.9), $\bar{c}_H(M) \sim O(v^2/f^2)$, $\bar{c}_{W,B}(M) \sim O(m_W^2/M^2)$, it follows that the correction to $\bar{c}_{W+B}$ from its RG evolution down to the scale μ is of order $\Delta\bar{c}_{W+B}/\bar{c}_{W+B}(M) \sim (g_*^2/16\pi^2)\log(M/\mu)$ as anticipated. Similarly, the insertion of $\bar{c}_{H\psi}$ into a loop of fermions, like in diagram (b) of Fig. 1, leads to a renormalization of $\bar{c}_W$ and $\bar{c}_B$:

$$\Delta\bar{c}_{W,B} \approx N_c\frac{\alpha_2}{4\pi}\log\left(\frac{\mu}{M}\right)\bar{c}_{H\psi}(M)\,, \tag{4.59}$$

where $N_c = 3$ is a color factor. In this case the RG correction is of order $(\lambda_\psi^2/16\pi^2)\log(M/\mu)$ compared to the UV value of the coefficients, as one can immediately verify by using the estimates (2.9).



Figure 1: One-loop diagrams relevant for the RG running of $\bar{c}_W$ and $\bar{c}_B$. Dashed, continuous and wiggly lines denote, respectively, a weak doublet H, a fermion and a vector field $V = W, B$. The symbol $\otimes$ denotes the insertion of the effective vertex from O_H (in diagram (a)) or $O_{H\psi}$ (in diagram (b)).



Figure 2: One-loop diagrams relevant for the RG running of $\bar{c}_T$. Dashed, continuous and wiggly lines denote, respectively, a weak doublet H, a fermion and a hypercharge field B. The symbol $\otimes$ denotes the insertion of the effective vertex from O_H (in diagram (a)) or $O_{H\psi}$ (in diagram (b)).

Loops of EW gauge fields give corrections which are suppressed by a weak loop factor $(g^2/16\pi^2)$, and the associated RG evolution is therefore generically small. An important exception is the case in which the Wilson coefficient has a value suppressed at the scale M. For example, if the dynamics behind the EW symmetry breaking is custodially invariant, then $\bar{c}_T(M) = 0$. The insertion of $\bar{c}_H$ into a loop of hypercharge gauge bosons, as in diagram (a) of Fig. 2, renormalizes $\bar{c}_T$ and gives

$$\bar{c}_T(\mu) = \frac{3}{2}\tan^2\theta_W \, \frac{\alpha_2}{4\pi} \log\left(\frac{\mu}{M}\right) \bar{c}_H(M) \,. \tag{4.60}$$

Compared to the naive estimate of Eq. (2.9), $\bar{c}_T(M) \sim O(v^2/f^2)$, valid in absence of custodial symmetry, the above correction is further suppressed by a factor $(g'^2/16\pi^2)\log(M/\mu)$. Although small, such a low-energy value of $\bar{c}_T$ has a strong impact on the EW precision tests performed at LEP [60]. [18] On the other hand, it is too small to be observable through a measurement of the Higgs couplings at the LHC. A similar renormalization of $\bar{c}_T$ also follows from loops of SM fermions through the insertion of $\bar{c}_{H\psi}$, as illustrated by diagram (b) of Fig. 2. The explicit calculation for the case of a composite right- and left-handed top quark was performed for example in Ref. [61]. Naively, the effect goes like

$$\Delta\bar{c}_T \approx N_c \frac{y_\psi^2}{16\pi^2} \log\left(\frac{\mu}{M}\right) \bar{c}_{H\psi}(M)\,, \tag{4.61}$$

and is of order $(y_\psi/g')^2(\lambda_\psi/g_*)^2$ compared to the one from loops of hypercharge. [19]

In general, although small, the RG evolution of the Wilson coefficients due to EW loops must be properly taken into account in order to precisely match the experimental results obtained at low energy with the theory predictions at high energy. This is even more true in the case of QCD loop corrections, which can be large and will affect the coefficients of the dimension-6 operators with quarks and gluon fields. [20] The effect of the running of the Wilson coefficients can be easily incorporated in programs for the automatic calculation of production cross sections and decay rates by using the effective Lagrangian (2.1) and identifying the coefficients appearing there as their values at the relevant low-energy scale. This is, in particular, how we proceeded to build `eHDECAY`, the modified version of `HDECAY` presented in Appendix E.

4.2 Decay rates at the loop level with the effective Lagrangian

In addition to the short-distance effects discussed above, which are parametrized in terms of the evolution of the coefficients of local operators, one-loop diagrams also lead to long-distance corrections to the observables under consideration. Specifically, while short-distance effects are related to the divergent terms, the long-distance contributions correspond to the finite parts and are defined in a given renormalization scheme. In general, the decay amplitude can be expanded as follows: [21]

---

[18] For example, $\bar{c}_T(m_Z) \sim 10^{-3}$ for $\bar{c}_H(M) \sim 0.1$.

[19] Notice that in case of a sizable fermion coupling λ_ψ, a numerically larger contribution to $\bar{c}_T$ comes from fermionic loops with *two* insertions of $\bar{c}_{H\psi}$. The corresponding diagram is quadratically divergent, so that it gives a threshold correction to $\bar{c}_T$ at the scale M, but does not contribute to its running. An explicit calculation can be found in Ref. [61] for the case of a composite top quark. Naively the effect is of order $\Delta\bar{c}_T \sim N_c(v/f)^2(\lambda_\psi/16\pi^2)(\lambda_\psi/g_*)^2$, and can be numerically large. For example, if both t_L and t_R couple with the same strength $\lambda_{t_L} = \lambda_{t_R} \sim \sqrt{g_* y_t}$ to the new dynamics, then it follows $\Delta\bar{c}_T \sim N_c(v/f)^2(y_t^2/16\pi^2)$.

[20] Notice that $g_s^2\bar{c}_g$ is not renormalized at one-loop by QCD corrections. This follows from the RG-invariance of the operator $(\beta(g_s)/g_s)G_{\mu\nu}G^{\mu\nu}$ which contributes to the trace of the energy-momentum tensor [62–64]. See also the recent discussion in Ref. [58].

$$A = A_0^{SM} + A_1^{SM} + \Delta A_0 + \Delta A_1 + \ldots \tag{4.62}$$

where A_0^{SM} (A_1^{SM}) is the tree-level (one-loop) SM amplitude, and ΔA_0 (ΔA_1) is the tree-level (one-loop) contribution from the dimension-6 operators of the effective Lagrangian in Eqs. (2.2)–(2.4). The dots denote higher-loop contributions as well as the corrections due to higher-order operators.

Let us consider for example the decay $h \to W^{(*)}W^*$. In this case the operators that can contribute at tree-level are O_H, O_W, O_{HW}, $O_{\psi W}$, $O'_{H\psi}$, as well as O_{Hud} in the case in which the off-shell W decays into a pair of quarks. Based on the naive estimates of Eq. (2.9) and according to the discussion of Section 3, we can quantify the various effects encoded by ΔA_0 as follows:

$$\begin{aligned}\frac{\Delta A_0}{A_0^{SM}}(W^{(*)}W^*) = {} & \hat{c}_H \times O\left(\frac{v^2}{f^2}\right) + \hat{c}_W \times O\left(\frac{E^2}{M^2}\right) + \hat{c}_{HW} \times O\left(\frac{E^2}{16\pi^2 f^2}\right) \\ & + \hat{c}_{Hud} \times O\left(\frac{v^2}{f^2}\,\frac{\lambda_u \lambda_d}{g_*^2}\right) + \hat{c}'_{H\psi} \times O\left(\frac{v^2}{f^2}\,\frac{\lambda_\psi^2}{g_*^2}\right) + \hat{c}_{\psi W} \times O\left(\frac{E m_\psi}{16\pi^2 f^2}\right) .\end{aligned} \tag{4.63}$$

Here $E = m_h$ is the relevant energy of the process and we have conveniently defined each of the $O(1)$ parameters $\hat{c}_i$ to be equal to $\bar{c}_i(m_h)$ divided by its naive estimate in Eq. (2.9):

---

[21] In the strict sense this equation is valid for the genuine EW corrections only, while for simplicity we include the (IR-divergent) virtual QED corrections to the SM amplitude in the same way. The corresponding real photon radiation contributions to the decay rates are treated in terms of a *linear* novel contribution to the Higgs coupling for the squared amplitude in order to obtain an infrared finite result. Pure QED corrections factorize as QCD corrections in general so that their amplitudes scale with the modified Higgs couplings. However, they cannot be separated from the genuine EW corrections in a simple way.

$$\hat{c}_i = \frac{f^2}{v^2}\bar{c}_i(m_h), \quad i = H, T, 6, \psi, \qquad \hat{c}_i = \frac{M^2}{m_W^2}\bar{c}_i(m_h), \quad i = W, B\,,$$
$$\hat{c}_i = \frac{16\pi^2 f^2}{m_W^2}\bar{c}_i(m_h), \quad i = HW, HB, \gamma, g, \psi W, \psi B, \psi G\,, \tag{4.64}$$
$$\hat{c}_i = \frac{g_*^2}{\lambda_\psi^2}\frac{f^2}{v^2}\bar{c}_i(m_h), \quad \hat{c}'_i = \frac{g_*^2}{\lambda_\psi^2}\frac{f^2}{v^2}\bar{c}'_i(m_h), \quad i = H\psi\,, \qquad \hat{c}_{Hud} = \frac{g_*^2}{\lambda_u\lambda_d}\frac{f^2}{v^2}\bar{c}_{Hud}(m_h)\,.$$

When the Higgs boson is pNGB, the two parameters $\hat{c}_g$ and $\hat{c}_\gamma$ are not of order one but are further suppressed by a factor $g^2_{\not{G}}/g_*^2$. From Eq. (4.63) one can see that the contribution of the dipole operators $O_{\psi W}$ is suppressed by (m_ψ/m_h) compared to that of O_{HW}, while that of O_{Hud} and $O'_{H\psi}$ is expected to be small given the existing constraints on the couplings λ_ψ (see the discussion in Section 2.1). The dominant NP contribution thus comes from the terms in the first line of Eq. (4.63), among which the one proportional to $\bar{c}_H$ is the leading effect for $g_* > g$. The 1-loop electroweak amplitude A_1^{SM} gives a contribution of order $A_1^{SM}/A_0^{SM} \sim (\alpha_2/4\pi)$. We thus see explicitly that ΔA_0 and A_1^{SM} encode the NLO corrections in the three expansion parameters which we are considering: $\alpha_2/4\pi$ (electroweak expansion), E^2/M^2 (derivative expansion) and v^2/f^2. The contribution due to 1-loop diagrams with one insertion of the effective vertices has not been computed yet, but we can easily estimate its size:

$$\frac{\Delta A_1}{A_0^{SM}}(W^{(*)}W^*) = \hat{c}_H \times O\!\left(\frac{v^2}{f^2}\frac{\alpha_2}{4\pi}\right) + \hat{c}_u \times O\!\left(\frac{v^2}{f^2}\frac{\alpha_2}{4\pi}\right) + \hat{c}_6 \times O\!\left(\frac{v^2}{f^2}\frac{\alpha_2}{4\pi}\right) + \dots \tag{4.65}$$

where the dots denote the subleading terms due to the other operators. The terms shown in Eq. (4.65) arise from the same 1-loop diagrams that give the SM amplitude A_1^{SM}, where each of the Higgs couplings gets shifted by $\bar{c}_H$, $\bar{c}_u$ and $\bar{c}_6$. By neglecting the unknown ΔA_1 one is omitting terms of order $(v^2/f^2)(\alpha_2/4\pi)$, that is, of the same size of the tree-level contribution due to the operator O_{HW}, see Eq. (4.63), since $E = m_h \approx m_W$. This latter contribution can be easily computed and it is included in the formula of the decay rate to WW (and similarly that of O_{HW} and O_{HB} to ZZ is also included) implemented by the program `eHDECAY` discussed in the Appendix (see Eq.(E.131) and also Eqs.(3.40), (E.138), (E.139)). The addition of the tree-level correction from O_{HW} is clearly the first step towards a full inclusion of the $O[(v^2/f^2)(\alpha_2/4\pi)]$ corrections, where the missing part will have to be computed from 1-loop diagrams featuring one insertion of O_H, O_u and O_6. It is worth noting that these diagrams, in general, contain logarithmic divergences which must be reabsorbed by a renormalization of the Wilson coefficients and contribute to their RG evolution as explained in the previous section. The finite part is the contribution to ΔA_1 which awaits to be computed.

By approximating the amplitude as $A \simeq A_0^{SM} + A_1^{SM} + \Delta A_0$ one obtains the following formula for the decay rate: [22]

$$\begin{aligned}\Gamma(W^{(*)}W^*) = \Gamma_0^{SM}(W^{(*)}W^*)\Bigg\{ 1 + \frac{2}{|A_0^{SM}|^2}\,\mathrm{Re}\big[\big(A_0^{SM}\big)^*\big(A_1^{SM} + \Delta A_0\big)\big] \\ + O\left(\left(\frac{v^2}{f^2}\right)^2, \left(\frac{\alpha_2}{4\pi}\frac{v^2}{f^2}\right), \left(\frac{\alpha_2}{4\pi}\right)^2\right)\Bigg\},\end{aligned} \tag{4.66}$$

where $\Gamma_0^{SM}(W^{(*)}W^*)$ denotes the tree-level SM decay rate. For simplicity, we have not shown terms involving powers of E^2/M^2 among the neglected contributions, since for $E = m_h \approx m_W$ one has $E^2/M^2 \lesssim v^2/f^2$ if $g_* \gtrsim g$. As mentioned, this formula incorporates the $O(v^2/f^2)$, $O(\alpha_2/4\pi)$ and $O(m_h^2/M^2)$ corrections (NLO in the perturbative expansion), and can be easily implemented in existing codes for the automatic computation of the decay rate. The inclusion of the $O(m_h^2/M^2)$ tree-level correction due to O_W is justified as long as $g_* < 4\pi$, since it is parametrically larger than the neglected $O[(v^2/f^2)(\alpha_2/4\pi)]$ terms by a factor $(16\pi^2/g_*^2)$. Notice that in the limit of large deviations of the Higgs couplings from their SM values, $(v/f)^2 \sim O(1)$, the neglected terms of $O[(v^2/f^2)(\alpha_2/4\pi)]$ become as important as those included through A_1^{SM}. In order words, a proper inclusion of the EW corrections in the limit $v \sim f$ requires a complete 1-loop calculation where each of the diagrams is rescaled by the appropriate coupling factor.

A similar discussion applies to the Higgs decay into a pair of fermions, $h \to \bar{\psi}\psi$. In this case only O_H and O_ψ ($\psi = u, d, l$) contribute at tree level,

$$\frac{\Delta A_0}{A_0^{SM}}(\bar{\psi}\psi) = \left(\frac{\hat{c}_H}{2} + \hat{c}_\psi\right) \times O\left(\frac{v^2}{f^2}\right), \tag{4.67}$$

while the one-loop EW diagrams featuring one effective vertex give a correction of order

$$\frac{\Delta A_1}{A_0^{SM}}(\bar{\psi}\psi) = \hat{c}_H \times O\left(\frac{v^2}{f^2}\frac{\alpha_2}{4\pi}\right) + \hat{c}_\psi \times O\left(\frac{v^2}{f^2}\frac{\alpha_2}{4\pi}\right) + \hat{c}_6 \times O\left(\frac{v^2}{f^2}\frac{\alpha_2}{4\pi}\right) + \ldots \tag{4.68}$$

---

[22] The same remark as in footnote 21 applies.

where the dots indicate the subleading terms due to the other operators. The calculation of ΔA_1 has not been performed yet, while the 1-loop EW corrections are known in the SM, A_1^{SM}. Their inclusion is thus possible as long as $(v/f) \ll 1$, so that the neglected terms in ΔA_1 are subleading. The case of QCD radiative corrections is different, since at leading order they factorize with respect to the expansion in the number of derivative and fields and can thus be resummed up to higher orders. In the case of the Higgs decay into a pair of quarks one can for example approximate $A \simeq A_0^{SM} + A_1^{SM} + \Delta A_0$ and obtain the following formula for the decay rate: [23]

$$\begin{aligned}\Gamma(\bar{q}q) = \Gamma_0^{SM}(\bar{q}q)\, \kappa^{QCD} \bigg\{ 1 &+ \frac{2}{|A_0^{SM}|^2} \,\mathrm{Re}\big[\big(A_0^{SM}\big)^* \big(A_1^{SM} + \Delta A_0\big)\big] \\ &+ O\left(\left(\frac{v^2}{f^2}\right)^2, \left(\frac{\alpha_2}{4\pi}\frac{v^2}{f^2}\right), \left(\frac{\alpha_2}{4\pi}\right)^2\right) \bigg\}\,,\end{aligned} \tag{4.69}$$

where $\Gamma_0^{SM}(\bar{q}q)$ is the SM tree-level rate and κ^{QCD} encodes the QCD corrections. This formula includes the leading $O(v^2/f^2)$, $O(\alpha_2/4\pi)$ and QCD corrections. Mixed electroweak and QCD corrections have been also included by assuming that they factorize, as the non-factorizable terms are known to be small. Compared to the decay rate into WW, Eq. (4.69) apparently does not include corrections of order m_h^2/M^2. While there is indeed no operator whose contribution starts at that order, such corrections can arise from subleading contributions to $\bar{c}_H$ and $\bar{c}_\psi$. For example, the tree-level exchange of heavy fermions can lead to a wave-function renormalization of the SM ones, which can be re-expressed in our notation as a contribution to $\bar{c}_\psi$ of order $\lambda_\psi^2 v^2/M^2$.

A similar resummation of the QCD corrections also works for the decay $h \to gg$. In this case the SM tree-level amplitude vanishes, $A_0^{SM} = 0$, while the leading contribution arises from the 1-loop exchange of top quarks. The two-loop EW corrections are known in the SM and give a correction of order $A_2^{SM}/A_1^{SM} \sim \alpha_2/4\pi$. Among the dimension-6 operators, only O_g contributes at tree-level,

$$\frac{\Delta A_0}{A_1^{SM}}(gg) = \hat{c}_g \times O\left(\frac{v^2}{f^2}\right)\,. \tag{4.70}$$

As discussed in Section 2 (see Eq. (2.10)), the above estimate is suppressed by an additional factor $(g_{\not{G}}^2/g_*^2)$ in the case of a NG Higgs boson. At the one-loop level one has

---

[23] The same remark as in footnote 21 applies.

$$\frac{\Delta A_1}{A_1^{SM}}(gg) = \left(\frac{\hat{c}_H}{2} + \hat{c}_u\right) \times O\left(\frac{v^2}{f^2}\right) + \hat{c}_{tG} \times O\left(\frac{v^2}{f^2}\,\frac{y_t^2}{16\pi^2}\right) . \tag{4.71}$$

Thus, the one-loop effect of O_H and O_u is expected to be as important as the tree-level one from O_g, and even larger if the Higgs is a NG boson, as discussed in Section 3.3. This is in fact not surprising, since $\bar{c}_g$ arises at the 1-loop level in minimally coupled theories, while $\bar{c}_H$ and $\bar{c}_u$ can be generated at tree level. The contribution from the dipole operator O_{tG} is suppressed by a factor $y_t^2/16\pi^2$ compared to that from O_H and O_u, as expected from the fact that $\bar{c}_{tG}$ is generated at the 1-loop level in minimally coupled theories. For this reason it will be neglected in the following. It should be noted that without a complete computation of the NLO EW corrections of order $(\alpha_2/4\pi)(v^2/f^2)$, the LHC data on Higgs physics are not sensitive to the range of values of $\bar{c}_{tG}$ expected using the naive estimate (2.9) with $(v/f)^2 \sim 0.1$. Furthermore, we stress that in order to distinguish the effect of O_{tG} from that of O_g, the $t\bar{t}h$ channel should be measured [49] (single top production in association with the Higgs could also provide complementary information [65]). Also in this case, there are no operators giving m_h^2/M^2 corrections, although these terms will in general appear as subleading contributions to $\bar{c}_g$, $\bar{c}_H$ and $\bar{c}_u$, as discussed above. It is well known that higher-order α_s corrections are large, so they must be included consistently in our perturbative expansion. This can be done easily in the approximation $m_h \ll 2m_t$, which is reasonably accurate for $m_h = 125\,\text{GeV}$. In such a limit one can integrate out the top quark and match its one-loop contribution to that of the local operator O_g. Then it trivially follows that the QCD corrections associated to the virtual exchange and real emissions of gluons and light quarks below the scale m_t factorize in the rate, the multiplicative factor being the same for both the top quark and New Physics terms. By approximating $A \simeq A_1^{SM} + A_2^{SM} + \Delta A_0 + \Delta A_1$, one arrives at the following formula for the $h \to gg$ decay rate:

$$\begin{split}\Gamma(gg) = \Gamma_1^{SM}(gg)\,\kappa_{soft}\,\Bigg\{ c_{eff}^2 + \frac{2\,c_{eff}}{|A_1^{SM}|^2}\,\text{Re}\big[\big(A_1^{SM}\big)^*\,\big(A_2^{SM} c_{eff} + \Delta A_0 + \Delta A_1\,c_{eff}\big)\big] \\ + O\Bigg(\left(\frac{v^2}{f^2}\right)^2, \left(\frac{\alpha_2}{4\pi}\,\frac{v^2}{f^2}\right), \left(\frac{\alpha_2}{4\pi}\right)^2\Bigg)\Bigg\},\end{split} \tag{4.72}$$

where $\Gamma_1^{SM}(gg)$ is the 1-loop SM decay width. The factor c_{eff} includes all the dependence on m_t and accounts for virtual QCD corrections to A_1^{SM} above that scale, while

κ_{soft} parametrizes the soft radiative effects. By using Eq. (4.72), the existing four-loop calculations of c_{eff} [66–69] and κ_{soft} [70–72] allow one to include the QCD corrections up to N^{3}LO.

The contributions to the decay $h \to \gamma\gamma$ follow a similar pattern as for $h \to gg$. At tree level:

$$\frac{\Delta A_0}{A_1^{SM}}(\gamma\gamma) = \hat{c}_\gamma \times O\left(\frac{v^2}{f^2}\right) . \tag{4.73}$$

At one loop:

$$\begin{aligned}\frac{\Delta A_1}{A_1^{SM}}(\gamma\gamma) = &\, \hat{c}_H \times O\left(\frac{v^2}{f^2}\right) + \hat{c}_u \times O\left(\frac{v^2}{f^2}\right) + \hat{c}_W \times O\left(\frac{m_W^2}{M^2}\right) \\ &+ \hat{c}_{HW} \times O\left(\frac{m_W^2}{16\pi^2 f^2}\right) + (\hat{c}_{tW} + \hat{c}_{tB}) \times O\left(\frac{v^2}{f^2}\frac{y_t^2}{16\pi^2}\right) .\end{aligned} \tag{4.74}$$

The 2-loop electroweak corrections have been computed in the SM and can be included for $(v^2/f^2) \ll 1$, so that unknown $O[(v^2/f^2)(\alpha_2/4\pi)]$ effects arising from 2-loop diagrams with one effective vertex are negligible. From Eq. (4.74) one can see that the 1-loop contribution due to O_{HW} is of the same order as such neglected terms. Furthermore, it is currently not known and thus not included in the formulas implemented in `eHDECAY`, see Appendix E. The 1-loop correction from O_W, on the contrary, is parametrically larger by a factor $(16\pi^2/g_*^2)$ and should be included for $g_* < 4\pi$. The easiest way to compute it is by rewriting O_W in terms of the other operators through the equations of motions [4], see Eq. (3.31). The result is finite and has been included in the formulas implemented in `eHDECAY`. The 1-loop correction due to the dipole operators is suppressed by a factor $y_t^2/16\pi^2$ and can be neglected. Approximating $A \simeq A_1^{SM} + A_2^{SM} + \Delta A_0 + \Delta A_1$ one finds:

$$\begin{aligned}\Gamma(\gamma\gamma) = \Gamma_1^{SM}(\gamma\gamma) \Bigg\{ 1 &+ \frac{2}{|A_1^{SM}|^2} \mathrm{Re}\left[\left(A_1^{SM}\right)^* \left(A_2^{SM} + \Delta A_0 + \Delta A_1\right)\right] \\ &+ O\left(\left(\frac{v^2}{f^2}\right)^2, \left(\frac{\alpha_2}{4\pi}\frac{v^2}{f^2}\right), \left(\frac{\alpha_2}{4\pi}\right)^2\right) \Bigg\},\end{aligned} \tag{4.75}$$

Finally, the estimate of the corrections to $h \to \gamma Z$ is the following:

$$\frac{\Delta A_0}{A_1^{SM}}(Z\gamma) = \hat{c}_\gamma \times O\left(\frac{v^2}{f^2}\right) + (\hat{c}_{HW} - \hat{c}_{HB}) \times O\left(\frac{v^2}{f^2}\right) , \tag{4.76}$$

$$\begin{aligned} \frac{\Delta A_1}{A_1^{SM}}(Z\gamma) = {} & \hat{c}_H \times O\left(\frac{v^2}{f^2}\right) + \hat{c}_u \times O\left(\frac{v^2}{f^2}\right) + \hat{c}_W \times O\left(\frac{m_W^2}{M^2}\right) + \hat{c}_{HW} \times O\left(\frac{m_W^2}{16\pi^2 f^2}\right) \\ & + \hat{c}_{tW} \times O\left(\frac{v^2}{f^2}\frac{y_t^2}{16\pi^2}\right) + \hat{c}_{tB} \times O\left(\frac{v^2}{f^2}\frac{y_t^2}{16\pi^2}\right) . \end{aligned} \tag{4.77}$$

In this case the 1-loop electroweak corrections are not known in the SM, so that the formula for the decay rate reads:

$$\Gamma(Z\gamma) = \Gamma_1^{SM}(Z\gamma)\left\{1 + \frac{2}{|A_1^{SM}|^2}\,\mathrm{Re}\big[\big(A_1^{SM}\big)^* (\Delta A_0 + \Delta A_1)\big] + O\left(\left(\frac{v^2}{f^2}\right)^2, \left(\frac{\alpha_2}{4\pi}\right)\right)\right\}, \tag{4.78}$$

where only the contributions from O_H, O_u and O_W should be retained in ΔA_1 for consistency.

Through the above discussion we sketched how the effective Lagrangian can be implemented beyond the tree level in the calculation of physical quantities. In the case of the Higgs partial decay widths, in particular, we have seen how the EW and QCD corrections can be included consistently with the expansion in the number of fields and derivatives. As a more concrete illustration of these considerations, we have written a modified version of the program `HDECAY`, which we dub `eHDECAY`, where the corrections from all the local operators of the effective Lagrangian (2.2) are included at NLO. A detailed description of the code is given in Appendix E, where more explicit formulas for each of the Higgs partial widths are provided.

5 Discussion

The discovery of a resonance with a mass around 125 GeV similar to the long-sought Standard Model Higgs boson brings the exploration of the electroweak symmetry breaking sector under quantitative scrutiny. The LHC experiments, together with those at the Tevatron, report the signal strengths, i.e. the product of the Higgs production cross section times its decay branching ratio, for various final state channels. The main task of the community is now to interpret these data and understand the implications for the theory of New Physics that is expected to lie beyond the weak scale.

The EW oblique parameters provide a bound on the scale of New Physics but do not give detailed information about the nature of the NP sector. In order to understand how the weak scale is stabilized at the quantum level, i.e. how the hierarchy problem is solved, one crucial question is whether EW symmetry breaking proceeds by weak or strong dynamics. The direct observation of new degrees of freedom would provide a straightforward answer. But a glimpse of New Physics can also be caught from a dedicated study of the Higgs boson itself, and in particular from a measurement of its couplings, if a departure from the SM predictions is ever observed. It is useful to parametrize the deviations from the SM by the effective Lagrangian of Eq. (2.1). By measuring its Wilson coefficients $\bar{c}_i$ one can infer what kind of UV theory completes the SM.

If the coupling strength of the Higgs boson to the NP sector is of the order of the SM weak couplings, $g_* \approx g$, then our power counting (2.9) shows that the coefficients of the operators that can be generated at tree-level, $O_H, O_{u,d,l}, O_W$ and O_B, are expected to be all of the same order, m_W^2/M^2, where M is the typical mass scale of the NP spectrum, unless some special selection rule suppresses some of them. It is instructive to examine the predictions of the archetypal example of weakly-coupled UV completions: the Minimal Supersymmetric Standard Model (MSSM). First, R-parity protects the EW oblique parameters from any tree-level contributions, hence $\bar{c}_W$ and $\bar{c}_B$ are of order $(m_W^2/M^2)(\alpha_2/4\pi)$ and thus small. Second, the couplings of the lightest Higgs boson to the massive gauge bosons are given by $c_V = \sin(\beta - \alpha)$, where α is the rotation angle to diagonalize the CP-even mass matrix and $\tan\beta$ is the ratio of the vacuum expectation values of the two neutral CP-even Higgs bosons. In the decoupling limit, $\alpha \to \beta - \pi/2$, one has $c_V = 1 + O(m_Z^4/m_H^4)$, where m_H is the mass of the heaviest CP-even scalar (for a general treatment of the decoupling limit see for example Ref. [73]). This means that at tree-level the deviations of the Higgs-gauge boson couplings are generated by dimension-8 operators [74], while $\bar{c}_H$ arises only through loop effects and is naively of order $(m_W^2/M^2)(\alpha_2/4\pi)$. At the same time, the couplings to up- and down-type quarks read, respectively,

$$\begin{aligned} c_u &= +\frac{\cos\alpha}{\sin\beta} = 1 + 2\,\frac{m_Z^2}{m_H^2}\cos^2\!\beta\cos 2\beta + O\!\left(\frac{m_Z^4}{m_H^4}\right) \\ c_d &= -\frac{\sin\alpha}{\cos\beta} = 1 - 2\,\frac{m_Z^2}{m_H^2}\sin^2\!\beta\cos 2\beta + O\!\left(\frac{m_Z^4}{m_H^4}\right)\,. \end{aligned} \tag{5.79}$$

For moderately large $\tan\beta$ this implies $\bar{c}_d \sim m_Z^2/m_H^2$, while $\bar{c}_u$ is further suppressed by a factor $\sim 1/\tan^2\beta$ (see for example Refs. [75, 76] and the recent discussion in Ref. [77]). A pattern with small values of $\bar{c}_H, \bar{c}_W$, $\bar{c}_B$ and $\bar{c}_u$ but with a $\sim 15\%$ enhancement of the Higgs coupling to down-type quarks due to $\bar{c}_d$, for example, would be indicative of the MSSM with large $\tan\beta$ and the additional Higgs bosons around 300 GeV. Generic two-Higgs doublet models lead to a similar pattern of couplings, while models where the Higgs boson mixes with a scalar that is singlet under the SM gauge group can generate $\bar{c}_H$ at the tree level. In the MSSM, loops of light stops or staus as well as charginos can also give sizable contributions to the effective couplings of the light Higgs boson to photons and gluons, with $\bar{c}_g$, $\bar{c}_\gamma$ satisfying the naive estimates (2.9). For example, loops of stops lead to $\bar{c}_g \sim (g_*^2/16\pi^2)(m_W^2/m_{\tilde{t}}^2)$, where $g_* = y_t$ or $A_t/m_{\tilde{t}}$.

This situation has to be contrasted with the case of strongly coupled theories. There, our power counting (2.9) singles out $\bar{c}_H$, $\bar{c}_{u,d}$ as the dominant Wilson coefficients ($\bar{c}_6$ controls only the Higgs self-interaction and measuring it at the LHC will be challenging), while $\bar{c}_W$ and $\bar{c}_B$ are suppressed by the ratio $(g/g_*)^2$. Furthermore, a composite Higgs boson can be naturally light if it is the pseudo Nambu-Goldstone boson associated to the dynamical breaking of a global symmetry of the strong dynamics. This implies that the coefficients $\bar{c}_g$ and $\bar{c}_\gamma$ will also be suppressed by a factor $(g_{\not{G}}/g_*)^2$, where $g_{\not{G}}$ is some weak spurion breaking the Goldstone symmetry. The modifications in the gluon-fusion production cross section and in the decay rate to photons are thus controlled by $\bar{c}_H$ and $\bar{c}_u$.

The harvest of data collected by the LHC certainly calls for a definite theoretical framework to describe the Higgs-like resonance and compute production and decay rates accurately in perturbation theory without restricting to the SM hypothesis. Effective Lagrangians are one of the tools at our disposal to achieve this goal. Elaborating on the operator classification of Ref. [4], we estimated the present bounds on the Wilson coefficients and provided accurate expressions for the Higgs decay rates including various effects that were previously omitted in the literature. Assuming that the observed Higgs-like resonance is a spin-0 and CP-even particle, we discussed two general formulations of the effective Lagrangian, one of which relies on the linear realization of $SU(2)_L \times U(1)_Y$ at high energies. One of the questions that can be addressed by considering these two parametrizations is whether the theory of New Physics flows to the SM in the infrared, that is, whether the Higgs-like resonance is part of an EW doublet. If all the Higgs signal strengths measured at the LHC converge towards the SM prediction, it would be a very suggestive indication that indeed the Higgs boson combines together with the longitudinal components of the W and Z to form an EW doublet, since any other alternative requires some tuning to fake the SM rates. On the other hand, the doublet nature of the Higgs boson would be less obvious to establish if the signal strengths exhibit deviations from their SM predictions. We have pointed out that, if the EWSB dynamics is custodially symmetric, it is not possible to test whether the Higgs boson is part of a doublet by means of single-Higgs processes alone. A direct proof can come only from processes with multi-Higgs bosons in the final states [54], which are however challenging to study at the LHC. Precisely establishing the CP nature of the Higgs boson is another question that also requires accurate computations. If there is little doubt that the observed resonance has a large CP-even component, the possibility of a small mixing with a CP-odd component remains alive, and dedicated analyses will have to be performed to bound the mixing angle between the two components. To this aim too, an effective Lagrangian including the CP-odd operators listed in Appendix C provides the theoretical framework where this question can be addressed quantitatively.

The absence so far of direct signals of New Physics at the LHC indicates that the road to unveil the origin of the electroweak symmetry breaking might be long and go through precision analyses rather than copious production of new particles. For such a task, the well established technology of effective field theories is the most powerful and general tool we have to analyze the Higgs data and put them into a coherent picture together with the existing experimental information without assuming the validity of the Standard Model. There is still time for the LHC to disprove this pessimistic eventuality by reporting the discovery of new light particles or large shifts in some of the Higgs couplings. It is clear, however, that if the New Physics continues to remain elusive, a precise investigation of the Higgs properties will become the most urgent programme in high-energy physics both for the experimental and the theoretical community.

Acknowledgments

We thank B. Gavela, A. De Rujula, J.R. Espinosa, A. Falkowski, E. Franco, L. Merlo, A. Pomarol, R. Rattazzi, F. Riva, L. Silvestrini, M. Trott for insightful discussions, and the participants of the LHC Higgs XS Working Group, in particular A. David, A. Denner, M. Dührssen, M. Grazzini, G. Passarino and G. Weiglein for discussions and comments. We also thank J.F. Kamenik for useful explanations on the results of Ref. [25]. This research has been partly supported by the European Commission under the ERC Advanced Grant 226371 MassTeV and the contract PITN-GA-2009-237920 UNILHC. C.G. is supported by the Spanish Ministry MICNN under contract FPA2010-17747. The work of R.C. was partly supported by the ERC Advanced Grant No. 267985 *Electroweak Symmetry Breaking, Flavour and Dark Matter: One Solution for Three Mysteries (DaMeSyFla).* M.M. is supported by the DFG SFB/TR9 Computational Particle Physics.

A SM Lagrangian: notations and conventions

In this Appendix, we collect the conventions used throughout this paper. The field content decomposes under $SU(3)_C \times SU(2)_L \times U(1)_Y$ as

$$H = (1,2,1/2),\ L_L^i = (1,2,-1/2),\ l_R^i = (1,1,-1), \tag{A.80}$$

$$q_L^i = (3,2,1/6),\ u_R^i = (3,1,2/3),\ d_R^i = (3,1,-1/3), \tag{A.81}$$

where the hypercharge is defined as $Y = Q - T_{3L}$, and $i = 1,2,3$ is a flavor index. The action of the gauge group is fully characterized by the conventions used to define the covariant derivative. For instance, for the left-handed quark doublet, we have

$$D_\mu q_L = \left(\partial_\mu - \frac{i}{2} g_S \lambda^a g_\mu^a - \frac{i}{2} g \sigma^i W_\mu^i - \frac{i}{6} g' B_\mu\right) q_L \tag{A.82}$$

where $\lambda^a, a = 1\ldots 8$, and $\sigma^i, i = 1\ldots 3$, are the usual Gell-Mann and Pauli matrices. Accordingly, the gauge-field strengths are defined as

$$G_{\mu\nu}^a = \partial_\mu g_\nu^a - \partial_\nu g_\mu^a + g_S f^{abc} g_\mu^b g_\nu^c\,, \tag{A.83}$$

where f^{abc} are the $SU(3)$ structure constants.

The Yukawa interactions of the up-type quarks involve the Higgs charge-conjugate doublet defined as

$$H^c = i\sigma^2 H^* . \tag{A.84}$$

The renormalizable Lagrangian of the SM thus reads:

$$\begin{aligned}\mathcal{L}_{SM} = &-\frac{1}{4}G^a_{\mu\nu}G^{a\,\mu\nu} - \frac{1}{4}W^i_{\mu\nu}W^{i\mu\nu} - \frac{1}{4}B_{\mu\nu}B^{\mu\nu} + (D^\mu H)^\dagger(D_\mu H)\\ &+ i\left(\bar{L}_L\gamma^\mu D_\mu L_L + \bar{l}_R\gamma^\mu D_\mu l_R + \bar{q}_L\gamma^\mu D_\mu q_L + \bar{u}_R\gamma^\mu D_\mu u_R + \bar{d}_R\gamma^\mu D_\mu d_R\right)\\ &+ \mu_H^2 H^\dagger H - \lambda(H^\dagger H)^2 + (y_u\,\bar{q}_L H^c u_R + y_d\,\bar{q}_L H d_R + y_l\,\bar{L}_L H l_R + h.c.)\end{aligned} \tag{A.85}$$

B Electroweak Chiral Lagrangian in non-unitary gauge

We report here the expression of the EW chiral Lagrangian valid in a generic gauge and in the most general case in which the $SU(2)_L \times U(1)_Y$ is non-linearly realized. For simplicity, we will restrict to the case in which the EWSB dynamics has a custodial invariance. The scalar h is assumed to be CP-even and a singlet of the custodial symmetry, and does not necessarily belong to an $SU(2)_L$ doublet. The Lagrangian can be expanded in terms with an increasing number of derivatives

$$\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{EWSB}\,, \qquad \mathcal{L}_{EWSB} = -V(h) + \mathcal{L}^{(2)} + \mathcal{L}^{(4)} + \dots \tag{B.86}$$

where $\mathcal{L}_0$ contains the kinetic terms of the $SU(3)_c \times SU(2)_L \times U(1)_Y$ gauge fields and of the SM fermions, $\mathcal{L}_{EWSB}$ describes the sector responsible for EWSB, and $V(h)$ is the potential for h [34]:

$$V(h) = \frac{1}{2}m_h^2 h^2 + c_3\,\frac{1}{6}\left(\frac{3m_h^2}{v}\right) h^3 + \dots \tag{B.87}$$

Under the request of $SU(2)_V$ custodial symmetry, the longitudinal W and Z polarizations correspond to the NG bosons of the global coset $SU(2)_L \times SU(2)_R/SU(2)_V$ and are described by the 2×2 matrix

$$\Sigma(x) = \exp\left(i\sigma^a\chi^a(x)/v\right)\,, \tag{B.88}$$

where σ^a are the Pauli matrices. $SU(2)_L \times U(1)_Y$ (local) transformations read as

$$\Sigma(x) \to U_L\Sigma(x)U_Y^\dagger\,, \qquad U_L = \exp(i\alpha_L^a\sigma^a)\,, \quad U_Y = \exp(i\alpha_Y\sigma^3) \tag{B.89}$$

and the covariant derivative is defined by

$$D_\mu\Sigma = \partial_\mu\Sigma - \frac{i\,g}{2}W^a_\mu\,\sigma^a\,\Sigma + \frac{i\,g'}{2}B_\mu\,\Sigma\,\sigma^3\,. \tag{B.90}$$

At the level of two derivatives one has [34]:

$$\begin{aligned}
\mathcal{L}^{(2)} = &\frac{1}{2}(\partial_\mu h)^2 + \frac{v^2}{4}\mathrm{Tr}\left(D_\mu\Sigma^\dagger D^\mu\Sigma\right)\left(1 + 2c_V\,\frac{h}{v} + \cdots\right) \\
&- \frac{v}{\sqrt{2}}\lambda^u_{ij}\,\left(\bar{u}^{(i)}_L, \bar{d}^{(i)}_L\right)\Sigma\left(u^{(i)}_R, 0\right)^T\left(1 + c_u\,\frac{h}{v} + \cdots\right) + h.c. \\
&- \frac{v}{\sqrt{2}}\lambda^d_{ij}\,\left(\bar{u}^{(i)}_L, \bar{d}^{(i)}_L\right)\Sigma\left(0, d^{(i)}_R\right)^T\left(1 + c_d\,\frac{h}{v} + \cdots\right) + h.c. \\
&- \frac{v}{\sqrt{2}}\lambda^l_{ij}\,\left(\bar{\nu}^{(i)}_L, \bar{l}^{(i)}_L\right)\Sigma\left(0, l^{(i)}_R\right)^T\left(1 + c_l\,\frac{h}{v} + \cdots\right) + h.c.
\end{aligned} \tag{B.91}$$

where the dots stand for terms with two or more Higgs fields and an implicit sum over flavor indices $i, j = 1, 2, 3$ has been understood. After rotating to the fermion mass eigenbasis and by choosing the unitary gauge $\Sigma(x) = 1$, the sum of (B.87) and (B.91) coincides with the first two lines of Eq. (3.46) with $c_W = c_Z = c_V$.

At the level of four derivatives, there are 6 independent bosonic operators [24] which affect cubic vertices with one h field: [25]

$$\begin{aligned}
\mathcal{L}^{(4)} = &\,c'_{WW}\,W^a_{\mu\nu}W^{\mu\nu\,a}\,\frac{h}{v} + c'_{WB}\,\mathrm{Tr}\big(\Sigma^\dagger\,W^a_{\mu\nu}\sigma^a\,\Sigma\,B_{\mu\nu}\sigma^3\big)\,\frac{h}{v} + c'_{BB}\,B_{\mu\nu}B^{\mu\nu}\,\frac{h}{v} \\
&+ \frac{c'_W}{m_W}\,D^\mu W^a_{\mu\nu}\,\mathrm{Tr}\Big(\Sigma^\dagger\sigma^a i\overleftrightarrow{D}_\nu\Sigma\Big)\,h - \frac{c'_B}{m_W}\,\partial^\mu B_{\mu\nu}\,\mathrm{Tr}\Big(\Sigma^\dagger i\overleftrightarrow{D}_\nu\Sigma\,\sigma^3\Big)\,h \\
&+ \frac{c_{gg}}{2}\,G^a_{\mu\nu}G^{a\,\mu\nu}\,\frac{h}{v} + \ldots
\end{aligned} \tag{B.93}$$

The dots stand for terms which have two or more h fields or do not lead to cubic vertices, see Refs. [52, 53] for the complete list of bosonic operators in $\mathcal{L}^{(4)}$. In the unitary gauge,

---

[24] The operator $O_2 = (v/m_W^2)\,\mathrm{Tr}\big[(D_\mu\Sigma)^\dagger(D_\nu\Sigma)\big]\,\partial^\mu\partial^\nu h$ that appeared in Eq. (B.85) of the first version of this paper is actually redundant and can be eliminated in terms of the operators O'_{WW} and O'_{WB}. For example, the shift $c'_{WW} \to c'_{WW} + c_2$ and $c'_{WB} \to c'_{WB} + \tan^2\theta_W\,c_2$ can be used to remove c_2.

[25] Another convenient basis, which can be more easily compared to Eq. (3.46), is one in which the first two operators of Eq. (B.93) are replaced by

$$W^a_{\mu\nu}\,\mathrm{Tr}\Big[\Sigma^\dagger\sigma^a i\overleftrightarrow{D}^\mu\Sigma\Big]\,\partial^\nu h\,, \qquad B_{\mu\nu}\,\mathrm{Tr}\Big[\Sigma^\dagger i\overleftrightarrow{D}^\mu\Sigma\sigma^3\Big]\,\partial^\nu h\,. \tag{B.92}$$

This is in fact the basis adopted in Ref. [52].

Eq. (B.93) coincides with the last three lines of Eq. (3.46). More specifically, the coefficients c_{WW}, c_{ZZ}, $c_{Z\gamma}$, $c_{\gamma\gamma}$ can be written as linear combinations of c'_{WW}, c'_{BB}, c'_{WB},

$$\begin{aligned}
c_{WW} &= 2\, c'_{WW} \\
c_{ZZ} &= 2(\cos^2\theta_W\, c'_{WW} - 2\sin\theta_W \cos\theta_W\, c'_{WB} + \sin^2\theta_W\, c'_{BB}) \\
c_{\gamma\gamma} &= 2(\sin^2\theta_W\, c'_{WW} + 2\sin\theta_W \cos\theta_W\, c'_{WB} + \cos^2\theta_W\, c'_{BB}) \\
c_{Z\gamma} &= 2(\sin\theta_W \cos\theta_W\, c'_{WW} + \cos 2\theta_W\, c'_{WB} - \sin\theta_W \cos\theta_W\, c'_{BB})\,,
\end{aligned} \tag{B.94}$$

while $c_{W\partial W}$, $c_{Z\partial Z}$ can be expressed in terms of c'_W, c'_B:

$$\begin{aligned}
c_{W\partial W} &= 4c'_W \\
c_{Z\partial Z} &= 4\, c'_W + 4\tan\theta_W\, c'_B \\
c_{Z\partial\gamma} &= 4\tan\theta_W\, c'_W - 4\, c'_B\,.
\end{aligned} \tag{B.95}$$

Notice that Eqs. (B.94) and (B.95) are directly implied by Eq. (3.53), which follows from custodial invariance. It is simple to verify that the identities (3.47) and (3.48) are satisfied by the couplings appearing on the left-hand sides of respectively Eq. (B.94) and (B.95).

The above discussion shows explicitly that every operator in Eq. (3.46) can be dressed up with NG bosons and made manifestly invariant under local $SU(2)_L \times U(1)_Y$ transformations. [26]

The part of Eq. (B.86) which does not depend on the Higgs field h coincides with the non-linear chiral Lagrangian for $SU(2)_L \times U(1)_Y$ [78], in the limit of exact custodial symmetry. This latter assumption can be relaxed by specifying the sources of explicit breaking of the custodial symmetry, *i.e.* its spurions, in terms of which one can construct additional operators formally invariant under $SU(2)_L \times U(1)_Y$ local transformations. For example, the list of operators that follows in the case in which custodial invariance is broken by a field with the EW quantum numbers of hypercharge has been recently discussed in Ref. [53]. Since the choice of quantum numbers of the spurions is model-dependent (and in fact the strongest effects are expected to arise from the breaking due to the top quark, rather than hypercharge), we do not report here any particular list of operators, and prefer to refer to the existing literature for further details.

---

[26] Notice that h is *invariant* under $SU(2)_L \times SU(2)_R$ (hence $SU(2)_L \times U(1)_Y$) transformations. In the case in which h belongs to an $SU(2)_L$ doublet H, this follows from the fact that h parametrizes the norm of the doublet: $H^\dagger H = (v+h)^2/2$.

C Relaxing the CP-even hypothesis

If one relaxes the hypothesis that h is CP-even, there are six extra dimension-6 operators that need to be added to the effective Lagrangian (2.2):

$$\begin{aligned}\Delta\mathcal{L}_{CP} = & \frac{i\tilde{c}_{HW}\, g}{m_W^2}\,(D^\mu H)^\dagger\sigma^i(D^\nu H)\tilde{W}^i_{\mu\nu} + \frac{i\tilde{c}_{HB}\, g'}{m_W^2}\,(D^\mu H)^\dagger(D^\nu H)\tilde{B}_{\mu\nu} \\ & + \frac{\tilde{c}_\gamma\, g'^2}{m_W^2}\,H^\dagger H B_{\mu\nu}\tilde{B}^{\mu\nu} + \frac{\tilde{c}_g\, g_S^2}{m_W^2}\,H^\dagger H G^a_{\mu\nu}\tilde{G}^{a\mu\nu} \\ & + \frac{\tilde{c}_{3W}\, g^3}{m_W^2}\,\epsilon^{ijk}W^{i\,\nu}_{\mu}W^{j\,\rho}_{\nu}\tilde{W}^{k\,\mu}_{\rho} + \frac{\tilde{c}_{3G}\, g_S^3}{m_W^2}\,f^{abc}G^{a\,\nu}_{\mu}G^{b\,\rho}_{\nu}\tilde{G}^{c\,\mu}_{\rho}\,,\end{aligned} \tag{C.96}$$

where the dual field strengths are defined as $\tilde{F}_{\mu\nu} = \frac{1}{2}\epsilon_{\mu\nu\rho\sigma}F^{\rho\sigma}$ for $F = W, B, G$ (ϵ is the totally antisymmetric tensor normalized to $\epsilon_{0123} = 1$). Furthermore, the coefficients of the operators involving fermions will be in general complex numbers.

In the case of the effective chiral Lagrangian with $SU(2)_L \times U(1)_Y$ non-linearly realized, there are four additional operators, to be added to those of Eq. (B.93), which can affect cubic vertices with one h field:

$$\begin{aligned}\Delta\mathcal{L}^{(4)}_{CP} = & \tilde{c}'_{WW}\,\tilde{W}^a_{\mu\nu}W^{\mu\nu\,a}\,\frac{h}{v} + \tilde{c}'_{WB}\,\mathrm{Tr}\Big[\Sigma^\dagger\,\tilde{W}^a_{\mu\nu}\sigma^a\,\Sigma\,B_{\mu\nu}\sigma^3\Big]\,\frac{h}{v} \\ & + \tilde{c}'_{BB}\,\tilde{B}_{\mu\nu}B^{\mu\nu}\,\frac{h}{v} + \frac{\tilde{c}_{gg}}{2}\,\tilde{G}^a_{\mu\nu}G^{a\mu\nu}\,\frac{h}{v}\,.\end{aligned} \tag{C.97}$$

In the unitary gauge, both Lagrangians $\Delta\mathcal{L}_{CP}$ and $\Delta\mathcal{L}^{(4)}_{CP}$ are matched onto:

$$\begin{aligned}\Delta\mathcal{L}^{(4)}_{CP} = & \left(\tilde{c}_{WW}\,W^+_{\mu\nu}\tilde{W}^{-\mu\nu} + \frac{\tilde{c}_{ZZ}}{2}\,Z_{\mu\nu}\tilde{Z}^{\mu\nu} + \tilde{c}_{Z\gamma}\,Z_{\mu\nu}\tilde{\gamma}^{\mu\nu} + \frac{\tilde{c}_{\gamma\gamma}}{2}\,\gamma_{\mu\nu}\tilde{\gamma}^{\mu\nu} + \frac{\tilde{c}_{gg}}{2}\,G^a_{\mu\nu}\tilde{G}^{a\mu\nu}\right)\frac{h}{v} \\ & + \left(\tilde{c}_{W\partial W}\left(W^-_\nu D_\mu\tilde{W}^{+\mu\nu} + h.c.\right) + \tilde{c}_{Z\partial Z}\,\tilde{Z}_\nu\partial_\mu Z^{\mu\nu} + \tilde{c}_{Z\partial\gamma}\,Z_\nu\partial_\mu\tilde{\gamma}^{\mu\nu}\right)\frac{h}{v} + \dots\end{aligned} \tag{C.98}$$

When the EW symmetry is linearly realized, the coefficients of Eq. (C.98) are related to the Wilson coefficients of Eq. (C.96) through the same relations reported in Table 1 with the simple exchange $c_i \to \tilde{c}_i$ (and with $c_W = c_B = 0$). In the non-linear case, $\tilde{c}_{W\partial W} = \tilde{c}_{Z\partial Z} = \tilde{c}_{Z\partial\gamma} = 0$ while $\tilde{c}_{WW}, \tilde{c}_{ZZ}, \tilde{c}_{\gamma\gamma}$ and $\tilde{c}_{Z\gamma}$ are given in terms of the Wilson coefficients of Eq. (C.97) by relations identical to the ones of Eq. (B.94) (with $c_i \to \tilde{c}_i$ and $c_2 = 0$).

Finally, it should also be noted that when the CP-invariance assumption in the Higgs sector is relaxed, the couplings $c_{u,d,l}$ are allowed to take some complex values.

D Current bounds on dimension-6 operators

In this Appendix we explain how we derived the bounds on the coefficients of the dimension-6 operators reported in Section 2.1. For a given observable we construct a likelihood for the coefficients $\bar{c}_i$ as follows:

$$L(\bar{c}_i) \propto \exp\left[-(O_{SM} + \delta O(\bar{c}_i) - O_{exp})^2/(2\,\Delta O_{exp}^2)\right] \,, \tag{D.99}$$

where $O_{exp} \pm \Delta O_{exp}$ is the experimental value of the observable, O_{SM} denotes its SM prediction and $\delta O(\bar{c}_i)$ is the correction due to the effective operators. If several observables constrain the same coefficients $\bar{c}_i$, the global likelihood is constructed by multiplying those of each observable. We include the theoretical uncertainty on the SM prediction by integrating over a nuisance parameter whose distribution is appropriately chosen. We then quote the bound on a given coefficient by marginalizing over the remaining ones.

Let us consider for example the bounds of Eqs. (2.14) and (2.15). To derive them we used the EW fit performed in Ref. [20] by the GFitter collaboration, and constructed a likelihood for the various coefficients by computing their contributions to the Z-pole observables. For the latter, we used the SM predictions and experimental inputs reported in Table 1 of Ref. [20], treating the uncertainties on the SM predictions as normally distributed. We performed two separate fits: one on the coefficients of the operators involving the light quarks (u, d, s), and one on those with charged leptons and heavy quarks (c, b). We thus neglected, for simplicity, the correlations between these two sets of coefficients. The relevant observables in the first fit are Γ_{tot}, σ_{had} and R_l. They depend on the Wilson coefficients only through the following linear combination:

$$\begin{aligned} l = &\left(\frac{1}{4} - \frac{1}{3}\sin^2\theta_W\right)(\bar{c}_{Hq1} + \bar{c}'_{Hq1}) + \left(-\frac{1}{4} + \frac{1}{6}\sin^2\theta_W\right)(\bar{c}_{Hq1} - \bar{c}'_{Hq1} + \bar{c}_{Hq2} - \bar{c}'_{Hq2}) \\ &+ \frac{1}{3}\sin^2\theta_W\,\bar{c}_{Hu} - \frac{1}{6}\sin^2\theta_W\,(\bar{c}_{Hd} + \bar{c}_{Hs}) \,, \end{aligned} \tag{D.100}$$

which with 95% probability is constrained to lie in the interval

$$-\,0.62 \times 10^{-3} < l < 1.2 \times 10^{-3} \,. \tag{D.101}$$

Although there are no further observables at the Z-pole which can resolve the degeneracy implied by this result, we thought it useful to report the limits that one obtains from

Eq. (D.101) by turning on one coefficient at the time. These are the bounds reported in Eq. (2.14).

The second fit, performed on the coefficients of the operators with leptons and heavy quarks, makes use of all the observables at the Z pole and counts 7 unknowns, specifically: $x_i = \{(\bar{c}_{Hq2} + \bar{c}'_{Hq2}), \bar{c}_{Hc}, (\bar{c}_{Hq3} - \bar{c}'_{Hq3}), \bar{c}_{Hb}, \bar{c}_{Hl}, (\bar{c}_{HL} + \bar{c}'_{HL}), (\bar{c}_{HL} - \bar{c}'_{HL})\}$. For simplicity we assume lepton universality, and thus take the coefficients $\bar{c}_{Hl}$, $\bar{c}_{HL}$, $\bar{c}'_{HL}$ to be the same for all the leptonic generations. In terms of the above variables, the result of the fit is summarized by their central values $\bar{x}_i$, standard deviations σ_i and by the correlation matrix ρ_{ij}:

$$\begin{aligned}
\bar{c}_{Hq2} + \bar{c}'_{Hq2} &= (-5.5 \pm 5.1) \times 10^{-3} \\
\bar{c}_{Hc} &= (4.3 \pm 11) \times 10^{-3} \\
\bar{c}_{Hq3} - \bar{c}'_{Hq3} &= (3.0 \pm 3.0) \times 10^{-3} \\
\bar{c}_{Hb} &= (-3.5 \pm 1.5) \times 10^{-2} \\
\bar{c}_{Hl} &= (9.6 \pm 8.6) \times 10^{-4} \\
\bar{c}_{HL} + \bar{c}'_{HL} &= (-6.4 \pm 15) \times 10^{-4} \\
\bar{c}_{HL} - \bar{c}'_{HL} &= (-1.3 \pm 0.75) \times 10^{-3}
\end{aligned} \tag{D.102}$$

$$\rho = \begin{pmatrix}
1.0 & -0.82 & 0.028 & 0.00096 & -0.063 & -0.13 & -0.093 \\
-0.82 & 1.0 & 0.0057 & 0.00019 & -0.013 & -0.026 & -0.019 \\
0.028 & 0.0057 & 1.0 & -0.87 & 0.031 & 0.063 & 0.046 \\
0.00096 & 0.00019 & -0.87 & 1.0 & 0.0011 & 0.0022 & 0.0016 \\
-0.063 & -0.013 & 0.031 & 0.0011 & 1.0 & -0.15 & -0.63 \\
-0.13 & -0.026 & 0.063 & 0.0022 & -0.15 & 1.0 & -0.22 \\
-0.093 & -0.019 & 0.046 & 0.0016 & -0.63 & -0.22 & 1.0
\end{pmatrix} \tag{D.103}$$

The limits of Eq. (2.15) have been obtained by making use of the above formulas and marginalizing over all the coefficients except the one on which the bound is reported.

For the limits of Eqs. (2.11) and (2.12) we have used the fit on S and T performed in Ref. [20], by marginalizing on one parameter to extract the bound on the other.

To derive Eq. (2.17) we have used the theoretical predictions of the EDM of the neutron and mercury given in Ref. [23] in terms of the dipole moments of the quarks (see Eqs. (2.12), (3.65) and (3.71) of Ref. [23]), and the experimental results for these observables given respectively in Ref. [79] and Ref. [80]. We included the theoretical errors by assuming that they are uniformly distributed within the stated intervals. Only two linear combinations of the coefficients $\bar{c}_i$ can be constrained in this way, since two are the observables at disposal:

$$\begin{aligned} l_1 &= -\frac{2m_d}{m_W^2}\left[\mathrm{Im}(\bar{c}_{dG}) + 1.3\,\mathrm{Im}(\bar{c}_{dB} - \bar{c}_{dW})\right] - \frac{m_u}{m_W^2}\left[\mathrm{Im}(\bar{c}_{uG}) - 0.64\,\mathrm{Im}(\bar{c}_{uB} + \bar{c}_{uW})\right] \\ l_2 &= -\frac{2m_u}{m_W^2}\,\mathrm{Im}(\bar{c}_{uG}) + \frac{2m_d}{m_W^2}\,\mathrm{Im}(\bar{c}_{dG})\,. \end{aligned} \tag{D.104}$$

Using $m_u = 2.3\,\mathrm{MeV}$ and $m_d = 4.8\,\mathrm{MeV}$ we obtain, with 95% probability:

$$\begin{aligned} -1.59\times 10^{-12}\,\mathrm{GeV}^{-1} < l_1 < 1.78\times 10^{-12}\,\mathrm{GeV}^{-1} \\ -1.82\times 10^{-12}\,\mathrm{GeV}^{-1} < l_2 < 1.37\times 10^{-12}\,\mathrm{GeV}^{-1}\,. \end{aligned} \tag{D.105}$$

From the above result, by turning on one coefficient at the time, one obtains the limits given in Eq. (2.17). The bound on $\mathrm{Im}(\bar{c}_{tG})$ of Eq. (2.18) has been similarly derived from the neutron and mercury EDMs by following Ref. [25] and making use of the formulas given there.

The limits of Eq. (2.22) have been obtained from the experimental measurements of the electron [29] and muon [27] anomalous magnetic moments and their SM predictions (taken respectively from Ref. [30] and Refs. [27,28]). In this case we have included the theoretical errors by assuming that they are normally distributed. All the remaining bounds reported in Section 2.1, namely those of Eqs. (2.19)-(2.21) and Eq. (2.23) have been obtained by simply translating into our notation the results given in the references quoted in the text.

E eHDECAY: a full implementation of the Higgs effective Lagrangian into HDECAY

The program `HDECAY` [56] was originally written for the automatic computation of the Higgs partial decay widths and branching ratios in the SM and in its Minimal Supersymmetric extension (MSSM). Its latest version 5.10 includes the possibility of specifying modified couplings for up-type quarks, down-type quarks, leptons and vector bosons in the parametrization of Eq. (3.46), as well as of including the effective couplings c_{gg}, $c_{\gamma\gamma}$ and $c_{Z\gamma}$. We present here a modified version of the program labelled `eHDECAY`. It is available at the following URL: `http://www-itp.particle.uni-karlsruhe.de/~maggie/eHDECAY/`. In addition to the features already present in `HDECAY`, the new program includes the effective couplings c_{WW}, c_{ZZ}, $c_{W\partial W}$ and $c_{Z\partial Z}$, and thus fully implements the non-linear Lagrangian (3.46). [27] In fact, similarly to `HDECAY` 5.10, it also includes the possibility of choosing different couplings of the Higgs boson to each of the up and down quark flavors and lepton flavors. In this sense the program does not assume neither custodial symmetry nor flavor alignment. As explained in the text, Eq. (3.46) describes a generic CP-even scalar h at NLO in the derivative expansion. If h forms an $SU(2)_L$ doublet together with the longitudinal polarizations of the W and the Z, the Lagrangian can be expanded as in Eq. (2.2) for $(v/f) \ll 1$; in this case the values of the Higgs couplings c_i are given in the second column of Table 1. The program `eHDECAY` provides an option in its input file where the user can switch from the non-linear parametrization of Eq. (3.46) to that of the SILH Lagrangian Eq. (2.2). The user can also choose to set the values of the Higgs couplings to those predicted at tree-level in the benchmark composite Higgs models MCHM4 and MCHM5, see the last two columns of Table 1.

Similarly to the original version of `HDECAY`, all the relevant QCD corrections are included. They generally factorize with respect to the expansion in the number of fields and derivatives of the effective Lagrangian, and can thus be straightforwardly included by making use of the existing SM computations. The inclusion of the EW corrections is less straightforward and can be currently done in a consistent way only in the framework of the Lagrangian (2.2) and up to higher orders in (v/f). Going beyond such approximations would require fully dedicated computations which at the moment are not available in the literature. In `eHDECAY` the user has the option to include the one-loop EW corrections to a given decay rate only if the parametrization of Eq. (2.2) has been chosen. The same EW scheme used by `HDECAY`, with G_F, m_W and m_Z taken as input parameters, is also adopted in `eHDECAY`. The sine of the Weinberg angle is defined as

$$\sin^2\theta_W = 1 - \frac{m_W^2}{m_Z^2}\,, \tag{E.106}$$

---

[27] Notice that the operator proportional to $c_{Z\partial\gamma}$ does not affect the decay $h \to Z\gamma$ as long as the photon is on-shell.

following the so-called on-shell scheme [81]. Derived quantities in this scheme are also the electromagnetic coupling and the weak coupling. To describe this latter we have conveniently defined the parameter α_2 in Eq. (3.43). The formulas implemented in the program are thus written in terms of only the input parameters or their derived quantities $\sin\theta_W$ and α_2. The only exception to this rule is given by the decay rates $\Gamma(h \to \gamma\gamma)$ and $\Gamma(h \to Z\gamma)$, where we use the experimental value of the electromagnetic coupling in the Thomson limit, $\alpha_{em}(q^2 = 0)$, in order to resum large logarithms for on-shell photons.

Below it follows a detailed discussion of how the New Physics corrections are incorporated for each of the Higgs decay modes. We report explicitly the formulas implemented in the code and their level of approximation in the perturbative expansion of the effective Lagrangian. In all the following expressions, as explained in the text, the coefficients of the dimension-6 operators of the SILH Lagrangian (2.2) and those of the derivative operators of Eq. (3.46) must be identified with their values at the relevant low-energy scale $\mu = m_h$.

E.1 Decays into quarks and leptons

Upon adopting the effective description of the non-linear Lagrangian (3.46) and working at leading order in the derivative expansion, the Higgs boson partial decay width into a pair of fermions is obtained by rescaling the tree-level SM value $\Gamma_0^{SM}(\psi\bar{\psi})$ by a factor c_ψ^2. The QCD corrections to the decay widths into quarks which are currently available for the SM case include fully massive NLO corrections near threshold [82] and massless $\mathcal{O}(\alpha_s^4)$ corrections far above threshold [83–85]. Also, large logarithms can be resummed through the running of the quark masses and of the strong coupling constant. All these QCD corrections factorize with respect to the tree-level amplitude and can therefore be incorporated as done in `HDECAY` for the SM case. The decay rate can be written as follows:

$$\Gamma(\bar{\psi}\psi)\big|_{NL} = c_\psi^2\, \Gamma_0^{SM}(\bar{\psi}\psi)\left[1 + \delta_\psi\, \kappa^{QCD}\right] , \tag{E.107}$$

where Γ_0^{SM} denotes the leading-order decay width, $\delta_\psi = 1(0)$ for $\psi =$ quark (lepton) and κ^{QCD} encodes the QCD corrections.[28] This is the formula implemented by `eHDECAY` in the case of the non-linear Lagrangian (3.46). It is valid up to corrections of $O(m_h^2/M^2)$ in the derivative expansion and of $O(\alpha_2/4\pi)$ from EW loops. These latter corrections are available in the SM but contrary to the QCD ones do not factorize. Their inclusion in the case of generic Higgs couplings thus requires a dedicated calculation, which is not available at present.

---

[28] There is one caveat, however. In the case of decays into strange, charm or bottom quarks there are two-loop diagrams which involve loops of top quarks coupling to the Higgs boson. They need a rescaling different from c_ψ^2. It has been correctly taken into account by the appropriate modification factor $c_\psi c_t$ ($\psi = c, s, b$).

In case of the SILH parametrization, where the deviations of the Higgs couplings from their SM values are assumed to be of $O(v^2/f^2)$ and small, the decay rate can be written as

$$\Gamma(\bar{\psi}\psi)\big|_{SILH} = \Gamma_0^{SM}(\bar{\psi}\psi)\left[1 - \bar{c}_H - 2\bar{c}_\psi + \frac{2}{|A_0^{SM}|^2}\mathrm{Re}\left(A_0^{*SM} A_{1,ew}^{SM}\right)\right]\left[1 + \delta_\psi\, \kappa^{QCD}\right] , \quad \text{(E.108)}$$

where A_0^{SM}, $A_{1,ew}^{SM}$ are, respectively, the tree-level and EW one-loop [86] amplitudes of the SM. In this case the one-loop EW corrections can be easily included if one neglects terms of $O[(\alpha_2/4\pi)(v/f)^2]$ (the same remark as in footnote 21 applies). In particular, mixed contributions up to $O[(\alpha_2/4\pi)(\alpha_s/4\pi)^4]$ have been included by assuming that the electroweak and QCD corrections factorize, as the non-factorizable contributions are small. From the viewpoint of the expansion in inverse powers of the NP scale, formula (E.108) includes corrections of order $O(v^2/f^2)$. It neglects terms of $O(v^4/f^4)$, $O[(\alpha_2/4\pi)(v/f)^2]$, $O[(\alpha_2/4\pi)^2]$.

E.2 Decay into gluons

Upon selecting the Lagrangian (3.46), the rate into two gluons is computed in `eHDECAY` by means of the following formula:

$$\begin{aligned}\Gamma(gg)\big|_{NL} = \frac{G_F\alpha_s^2 m_h^3}{4\sqrt{2}\pi^3}\Bigg[&\left|\sum_{q=t,b,c}\frac{c_q}{3}\,A_{1/2}\left(\tau_q\right)\right|^2 c_{eff}^2\,\kappa_{soft} \\ &+ 2\,\mathrm{Re}\left(\sum_{q=t,b,c}\frac{c_q}{3}\,A_{1/2}^*\left(\tau_q\right)\frac{2\pi c_{gg}}{\alpha_s}\right) c_{eff}\,\kappa_{soft} + \left|\frac{2\pi c_{gg}}{\alpha_s}\right|^2 \kappa_{soft} \qquad \text{(E.109)} \\ &+ \frac{1}{9}\sum_{q,q'=t,b} c_q\, A_{1/2}^*\left(\tau_q\right) c_{q'}\, A_{1/2}\left(\tau_{q'}\right)\kappa^{NLO}(\tau_q,\tau_{q'})\Bigg] ,\end{aligned}$$

where $\tau_q = 4m_q^2/m_h^2$ and the loop function is defined as

$$A_{1/2}\left(\tau\right) = \frac{3}{2}\tau\left[1 + \left(1-\tau\right) f\left(\tau\right)\right] , \qquad \text{(E.110)}$$

with

$$f(\tau) = \begin{cases} \arcsin^2 \dfrac{1}{\sqrt{\tau}} & \tau \geq 1 \\ -\dfrac{1}{4}\left[\ln \dfrac{1+\sqrt{1-\tau}}{1-\sqrt{1-\tau}} - i\pi\right]^2 & \tau < 1\,. \end{cases} \tag{E.111}$$

The first term corresponds to the one-loop contribution from the top, bottom and charm quarks, whose couplings to the Higgs boson are modified with respect to their SM values. The second and third terms encode the effect of the derivative interaction between the Higgs boson and two gluons generated by New Physics. Naively $c_{gg} \approx (\alpha_s/4\pi)(g_*^2 v^2/M^2)$, so that the correction from the effective interaction can be as important as the one from the top quark if $(g_*^2 v^2/M^2) \approx 1$. No expansion is thus possible in c_{gg} in the general case.

The QCD corrections have been included up to N^{3}LO in Eq. (E.109) in the limit of heavy loop particle masses, similarly to what is done in `HDECAY` for the SM. In this limit the effect of soft radiation factorizes and is encoded by the coefficient κ_{soft}. The coefficient c_{eff}, instead, takes into account the correction from the exchange of hard gluons and quarks with virtuality $q^2 \gg m_t^2$. More in detail, for $m_h \ll 2m_t$, one can integrate out the top quark and obtain the following five-flavour effective Lagrangian

$$\mathcal{L}_{\text{eff}} = -2^{1/4} G_F^{1/2} C_1\, G^0_{a\mu\nu} G^{0\mu\nu}_a h\,, \tag{E.112}$$

where bare fields are labeled by the superscript 0. The renormalized coefficient function C_1 encodes the dependence on the top quark mass m_t. The coefficients κ_{soft} and c_{eff} are thus defined as

$$\begin{aligned} \kappa_{soft} &= \frac{\pi}{2m_h^4}\,\mathrm{Im}\,\Pi^{GG}(q^2 = m_h^2) \\ c_{eff} &= -\frac{12\pi\, C_1}{\alpha_s^{(5)}(m_h)}\,, \end{aligned} \tag{E.113}$$

where $\Pi^{GG}(q^2)$ is the vacuum polarization induced by the gluon operator. The N^{3}LO expression of the coefficient function C_1 [66–69] in the on-shell scheme and that of $\mathrm{Im}\,\Pi^{GG}$ can be found in Ref. [72]. At NLO the expressions for κ_{soft} and c_{eff} take the well-known form

$$\kappa^{NLO}_{soft} = 1 + \frac{\alpha_s^{NLO}}{\pi}\left(\frac{73}{4} - \frac{7}{6}N_F\right)\,, \qquad c^{NLO}_{eff} = 1 + \frac{\alpha_s^{NLO}}{\pi}\frac{11}{4}\,, \tag{E.114}$$

where here α_s^{NLO} is evaluated at the scale m_h and computed at NLO for $N_F = 5$ active flavours, while it has been consistently computed up to N^{3}LO for the inclusion in `eHDECAY`. The last line in Eq. (E.109) contains the additional mass effects at NLO QCD [71] in the top and bottom loops, encoded in $\kappa^{NLO}(\tau_q, \tau_{q'})$, which have been explicitly implemented in `HDECAY` and taken over in `eHDECAY`. While the mass effects for the top quark loops play only a minor role, below the percent level, for the bottom loop contribution the mass effects for a 125 GeV Higgs boson amount to about 8% relative to the approximate NLO result. Hence, formula (E.109) includes the QCD corrections at N^{3}LO (i.e. at $O(\alpha_s^5)$ in the decay rate), and neglects terms of $O(m_h^2/M^2)$ in the derivative expansion.

When the SILH Lagrangian (2.2) is selected, on the other hand, `eHDECAY` computes the decay rate into gluons by means of the following approximate formula:

$$
\begin{aligned}
\Gamma(gg)\big|_{SILH} = \frac{G_F \alpha_s^2 m_h^3}{4\sqrt{2}\pi^3} \Bigg[& \frac{1}{9} \sum_{q,q'=t,b,c} (1 - \bar{c}_H - \bar{c}_q - \bar{c}_{q'})\, A^*_{1/2}(\tau_{q'})\, A_{1/2}(\tau_q)\; c^2_{eff}\, \kappa_{soft} \\
& + 2\,\mathrm{Re}\left(\sum_{q=t,b,c} \frac{1}{3} A^*_{1/2}(\tau_q) \frac{16\pi\, \bar{c}_g}{\alpha_2} \right) c_{eff}\, \kappa_{soft} \\
& + \left| \sum_{q=t,b,c} \frac{1}{3} A_{1/2}(\tau_q) \right|^2 c^2_{eff}\, \kappa_{ew}\, \kappa_{soft} \\
& + \frac{1}{9} \sum_{q,q'=t,b} (1 - \bar{c}_H - \bar{c}_q - \bar{c}_{q'}) A^*_{1/2}(\tau_q)\, A_{1/2}(\tau_{q'})\, \kappa^{NLO}(\tau_q, \tau_{q'}) \Bigg] .
\end{aligned}
\tag{E.115}
$$

The last line contains the mass effects at NLO QCD for the top and bottom quark loops. The NLO electroweak corrections [87, 88] are included through the coefficient κ_{ew} and by neglecting terms of $O[(\alpha_2/4\pi)(v^2/f^2)]$. The above formula thus includes the leading $O(v^2/f^2)$ corrections, as well as mixed $O[(\alpha_s/4\pi)^5(\alpha_2/4\pi)]$ ones. Indeed, we assume factorisation of the QCD and EW corrections. Since QCD corrections are dominated by soft gluon radiation, in which QCD and EW effects completely factorize, this is a good approximation[29]. It neglects terms of $O[(\alpha_2/4\pi)^2]$ and $O(v^4/f^4)$.

---

[29] Bottom loops contribute $O(10\%)$ to the SM decay rate and are well approximated by an effective coupling at the 10%-level thus leading to negligible non-factorizing contributions at the per-cent level.

E.3 Decay into photons

In the SM the decay of the Higgs boson into a pair of photons is mediated by W and heavy fermion loops. According to the chiral Lagrangian (3.46), these two contributions to the total amplitude are rescaled, respectively, by the parameters c_V and c_ψ. Similarly to $h \to gg$, the contact interaction proportional to $c_{\gamma\gamma}$ can also contribute significantly. With $c_{\gamma\gamma} \approx (\alpha_{em}/4\pi)(g_*^2 v^2/M^2)$, the contribution due to the effective interaction becomes comparable to the loop induced contributions if $(g_*^2 v^2/M^2) \approx 1$. The partial width for a Higgs boson decaying into two photons implemented in `eHDECAY` in the framework of the non-linear Lagrangian is thus given by

$$\begin{aligned}\Gamma(\gamma\gamma)\big|_{NL} = \frac{G_F \alpha_{em}^2 m_h^3}{128\sqrt{2}\pi^3}\Bigg| & \sum_{q=t,b,c} \frac{4}{3} c_q \, 3Q_q^2 \, A_{1/2}^{NLO}(\tau_q) + \frac{4}{3} c_\tau Q_\tau^2 A_{1/2}(\tau_\tau) \\ & + c_V A_1(\tau_W) + \frac{4\pi}{\alpha_{em}} c_{\gamma\gamma}\Bigg|^2 ,\end{aligned} \tag{E.116}$$

which is approximate at leading order in the derivative expansion, i.e. it neglects terms of $O(m_h^2/M^2)$. Note that α_{em} is the electromagnetic coupling in the Thomson limit, in order to resum large logarithms for on-shell photons. We have defined $\tau_i = 4m_i^2/m_h^2$ $(i = q, \tau, W)$, and the form factor

$$A_1(\tau) = -\left[2 + 3\tau + 3\tau(2-\tau) f(\tau)\right] , \tag{E.117}$$

while $A_{1/2}(\tau)$ is given in Eq. (E.110). The top, bottom and charm quark loops receive NLO QCD corrections, while the effective contact interaction does not. The NLO QCD corrected quark form factor is denoted in Eq. (E.116) by

$$A_{1/2}^{NLO}(\tau_q) = A_{1/2}(\tau_q)(1 + \kappa_{QCD}) , \tag{E.118}$$

where κ_{QCD} encodes the $O(\alpha_s/4\pi)$ QCD corrections [89].

In the case of the SILH parametrization, the EW corrections have been incorporated as well. It is useful to define the SM amplitude at LO and NLO QCD level as

$$A_X^{SM}(\gamma\gamma) = \sum_{q=t,b,c} \frac{4}{3} 3Q_q^2 \, A_{1/2}^X(\tau_q) + \frac{4}{3} Q_\tau^2 \, A_{1/2}(\tau_\tau) + A_1(\tau_W) \, , \qquad X = LO, NLO \, , \tag{E.119}$$

where $A_{LO}^{SM} \equiv A^{SM}$, and the deviation from the SM amplitude as

$$\begin{aligned}\Delta A(\gamma\gamma) = &- \sum_{q=t,b,c} \frac{4}{3}\left(\frac{\bar{c}_H}{2} + \bar{c}_q\right) 3Q_q^2\, A_{1/2}^{NLO}\left(\tau_q\right) - \left(\frac{\bar{c}_H}{2} + \bar{c}_\tau\right)\frac{4}{3} Q_\tau^2\, A_{1/2}\left(\tau_\tau\right) \\ &- \left(\frac{\bar{c}_H}{2} - 2\bar{c}_W\right) A_1\left(\tau_W\right) .\end{aligned} \tag{E.120}$$

The decay width implemented in `eHDECAY` in the SILH case is thus the following

$$\begin{aligned}\Gamma(\gamma\gamma)\big|_{SILH} = \frac{G_F \alpha_{em}^2 m_h^3}{128\sqrt{2}\pi^3} \Bigg\{ &|A_{NLO}^{SM}(\gamma\gamma)|^2 + 2\,\mathrm{Re}\Big(A_{LO}^{SM*}(\gamma\gamma)\, A_{ew}^{SM}(\gamma\gamma)\Big) \\ &+ 2\,\mathrm{Re}\left[A_{NLO}^{SM*}(\gamma\gamma)\left(\Delta A(\gamma\gamma) + \frac{32\pi \sin^2\theta_W\, \bar{c}_\gamma}{\alpha_{em}}\right)\right]\Bigg\} ,\end{aligned} \tag{E.121}$$

where $A_{ew}^{SM}(\gamma\gamma)$ denotes the SM amplitude which comprises the NLO electroweak corrections [87, 90]. Equation (E.121) includes the leading $O(v^2/f^2)$ and $O(m_h^2/M^2)$ corrections, while it neglects terms of order $O(v^4/f^4)$. The electroweak corrections are implemented up to NLO, neglecting corrections of $O[(\alpha_2/4\pi)(v^2/f^2)]$ and of $O[(\alpha_2/4\pi)^2]$. Finally, the QCD corrections are included up to NLO, and mixed terms of $O[(\alpha_2/4\pi)(\alpha_s/4\pi)]$ are neglected.

E.4 Decay into $Z\gamma$

In the SM the Higgs boson decay into a Z boson and a photon is mediated by a W boson and heavy fermion loops. Adopting the parametrization of the non-linear Lagrangian, the correction from the effective interaction due to the coupling $c_{Z\gamma}$ has to be considered, too, and it can become as important as the loop contributions for $(g_*^2 v^2/M^2) \approx 1$. The decay width is therefore given by (here also $\alpha_{em} \equiv \alpha_{em}(0)$):

$$\begin{aligned}\Gamma(Z\gamma)\big|_{NL} = &\frac{G_F^2 \alpha_{em} m_W^2 m_h^3}{64\pi^4}\left(1 - \frac{m_Z^2}{m_h^2}\right)^3 \\ &\times \left|\sum_\psi \frac{c_\psi N_c Q_\psi \hat{v}_\psi}{\cos\theta_W}\, A_{1/2}^{Z\gamma}\left(\tau_\psi, \lambda_\psi\right) + c_V\, A_1^{Z\gamma}\left(\tau_W, \lambda_W\right) - \frac{4\pi}{\sqrt{\alpha_{em}\alpha_2}}\, c_{Z\gamma}\right|^2 ,\end{aligned} \tag{E.122}$$

with $\tau_i = 4m_i^2/m_h^2$, $\lambda_i = 4m_i^2/m_Z^2$, $\hat{v}_\psi = 2I_\psi^3 - 4Q_\psi \sin^2\theta_W$ $(\psi = t, b, c, \tau)$ and the form factors are defined by

$$\begin{aligned} A_{1/2}^{Z\gamma}(\tau,\lambda) &= \left[I_1(\tau,\lambda) - I_2(\tau,\lambda)\right] , \\ A_1^{Z\gamma}(\tau,\lambda) &= \cos\theta_W \Big\{ 4\big(3 - \tan^2\theta_W\big) I_2\big(\tau,\lambda\big) \\ &\qquad + \Big[\Big(1 + \frac{2}{\tau}\Big)\tan^2\theta_W - \Big(5 + \frac{2}{\tau}\Big)\Big] I_1\big(\tau,\lambda\big)\Big\} . \end{aligned} \tag{E.123}$$

The functions I_1 and I_2 can be cast into the form

$$\begin{aligned} I_1(\tau,\lambda) &= \frac{\tau\lambda}{2(\tau-\lambda)} + \frac{\tau^2\lambda^2}{2(\tau-\lambda)^2}\left[f(\tau) - f(\lambda)\right] + \frac{\tau^2\lambda}{(\tau-\lambda)^2}\left[g(\tau) - g(\lambda)\right] \\ I_2(\tau,\lambda) &= -\frac{\tau\lambda}{2(\tau-\lambda)}\left[f(\tau) - f(\lambda)\right] , \end{aligned} \tag{E.124}$$

where $f(\tau)$ is defined in Eq. (E.111) and $g(\tau)$ reads

$$g(\tau) = \begin{cases} \sqrt{\tau - 1}\,\arcsin\dfrac{1}{\sqrt{\tau}} & \tau \geq 1 \\ \dfrac{\sqrt{1-\tau}}{2}\left[\ln\dfrac{1+\sqrt{1-\tau}}{1-\sqrt{1-\tau}} - i\pi\right] & \tau < 1 . \end{cases} \tag{E.125}$$

The QCD radiative corrections [91] are small and thus have been neglected, while the NLO EW corrections are unknown. Finally, Eq. (E.122) neglects terms of $O(m_h^2/M^2)$, which are of higher-order in the derivative expansion.

In the SILH parametrization the decay width is computed by `eHDECAY` according to the formula

$$\begin{aligned} \Gamma(Z\gamma)\big|_{SILH} &= \frac{G_F^2 \alpha_{em} m_W^2 m_h^3}{64\pi^4}\left(1 - \frac{m_Z^2}{m_h^2}\right)^3 \\ &\quad \times \Bigg\{ \left|A^{SM}(Z\gamma)\right|^2 + 2\,\mathrm{Re}\big(A^{SM*}(Z\gamma)\,\Delta A(Z\gamma)\big) \\ &\qquad + 2\,\mathrm{Re}\Big[-\frac{4\pi\tan\theta_W}{\sqrt{\alpha_{em}\alpha_2}}(\bar{c}_{HB} - \bar{c}_{HW} - 8\bar{c}_\gamma \sin^2\theta_W)\,A^{SM*}(Z\gamma)\Big]\Bigg\} , \end{aligned} \tag{E.126}$$

where we have defined the LO SM amplitude

$$A^{SM}(Z\gamma) = \sum_\psi \frac{N_c Q_\psi \hat{v}_\psi}{\cos\theta_W} A_{1/2}^{Z\gamma}(\tau_\psi, \lambda_\psi) + A_1^{Z\gamma}(\tau_W, \lambda_W) \tag{E.127}$$

and the deviation from the SM amplitude

$$\Delta A(Z\gamma) = -\sum_{\psi} \left(\frac{\bar{c}_H}{2} + \bar{c}_\psi\right) \frac{N_c Q_\psi \hat{v}_\psi}{\cos\theta_W} \, A^{Z\gamma}_{1/2}\left(\tau_\psi, \lambda_\psi\right) - \left(\frac{\bar{c}_H}{2} - 2\bar{c}_W\right) \, A^{Z\gamma}_{1}\left(\tau_W, \lambda_W\right) \ . \tag{E.128}$$

Equation (E.126) includes corrections of $O(v^2/f^2)$ and $O(m_h^2/M^2)$. The EW corrections are unknown, and small QCD radiative corrections have been neglected.

E.5 Decays into WW and ZZ bosons

The Higgs boson decay into a pair of massive vector bosons is important not only above the threshold, but also below. For example, in the SM with $m_h = 125$ GeV the branching ratio of $h \to WW$ is about 20%. In `HDECAY` various options are present to compute the partial decay widths with on-shell or off-shell bosons, controlled by the ON-SH-WZ input parameter. In `eHDECAY` we have implemented the case ON-SH-WZ=0, which includes the double off-shell decays $h \to W^*W^*, Z^*Z^*$. For this case, which is obviously the most complete as it takes into account both on-shell and off-shell contributions, the partial decay width $h \to V^*V^*$ ($V = W, Z$) can be written in the following compact form:

$$\Gamma(V^*V^*) = \frac{1}{\pi^2} \int_0^{m_h^2} \frac{dQ_1^2 \, m_V \Gamma_V}{\left(Q_1^2 - m_V^2\right)^2 + m_V^2 \Gamma_V^2} \int_0^{(m_h - Q_1)^2} \frac{dQ_2^2 \, m_V \Gamma_V}{\left(Q_2^2 - m_V^2\right)^2 + m_V^2 \Gamma_V^2} \, \Gamma(VV) \, , \tag{E.129}$$

where Q_1^2, Q_2^2 are the squared invariant masses of the virtual gauge bosons and m_V and Γ_V their masses and total decay widths. In the parametrization of Eq. (3.46), by defining

$$a_{VV} = c_{VV} \, \frac{m_h^2}{m_V^2} \, , \qquad a_{V\partial V} = \frac{c_{V\partial V}}{2} \, \frac{m_h^2}{m_V^2} \, , \tag{E.130}$$

the squared matrix element $\Gamma(VV)$ reads

$$\begin{aligned} \Gamma(VV)\big|_{NL} = \ \Gamma^{SM}(VV) \times \Bigg\{ & c_V^2 - 2c_V \left[\frac{a_{VV}}{2} \left(1 - \frac{Q_1^2 + Q_2^2}{m_h^2}\right) + a_{V\partial V} \frac{Q_1^2 + Q_2^2}{m_h^2} \right] \\ & + c_V a_{VV} \, \frac{\lambda\left(Q_1^2, Q_2^2, m_h^2\right) \ \left(1 - (Q_1^2 + Q_2^2)/m_h^2\right)}{\lambda\left(Q_1^2, Q_2^2, m_h^2\right) + 12 \, Q_1^2 Q_2^2 / m_h^4} \Bigg\} \, , \end{aligned} \tag{E.131}$$

with [92]

$$\Gamma^{SM}(VV) = \frac{\delta_V G_F m_h^3}{16\sqrt{2}\pi} \sqrt{\lambda\left(Q_1^2, Q_2^2, m_h^2\right)} \left(\lambda\left(Q_1^2, Q_2^2, m_h^2\right) + \frac{12 Q_1^2 Q_2^2}{m_h^4}\right) \ , \tag{E.132}$$

where $\delta_V = 2(1)$ for $V = W(Z)$ and $\lambda(x,y,z) \equiv (1 - x/z - y/z)^2 - 4xy/z^2$. The second and third term in Eq. (E.131) represent the interference between the tree-level contribution and the one from the derivative operators. They are of order $O(m_h^2/M^2)$, hence next-to-leading in the chiral expansion compared to the tree-level contribution; we have consistently neglected terms quadratic in a_{VV} and $a_{V\partial V}$, since they are of $O(m_h^4/M^4)$, which is beyond the accuracy of the effective Lagrangian (3.46).

In the SILH parametrization the squared matrix element $\Gamma(VV)$ implemented in `eHDECAY` reads

$$\Gamma(VV)\big|_{SILH} = \Gamma^{SILH}(VV) + \Gamma^{SM}(VV)\frac{2}{|A_0^{SM}|^2}\,\mathrm{Re}\left(A_0^{*SM} A_{ew}^{SM}\right)\ , \tag{E.133}$$

where A_0^{SM} denotes the SM LO amplitude and A_{ew}^{SM} is the SM amplitude which comprises the NLO EW corrections [93] (the same remark as in footnote 21 applies). Furthermore,

$$\begin{aligned}\Gamma^{SILH}(VV) = \ \Gamma^{SM}(VV) \times \Bigg\{ & 1 - \bar{c}_H - 2\left[\frac{\bar{a}_{VV}}{2}\left(1 - \frac{Q_1^2 + Q_2^2}{m_h^2}\right) + \bar{a}_{V\partial V}\frac{Q_1^2 + Q_2^2}{m_h^2}\right] \\ & + \bar{a}_{VV}\,\frac{\lambda\left(Q_1^2, Q_2^2, m_h^2\right)\,\left(1 - (Q_1^2 + Q_2^2)/m_h^2\right)}{\lambda\left(Q_1^2, Q_2^2, m_h^2\right) + 12\,Q_1^2 Q_2^2/m_h^4}\Bigg\}\,,\end{aligned} \tag{E.134}$$

where we have defined,

$$\begin{aligned}\bar{a}_{WW} &= -2\,\frac{m_h^2}{m_W^2}\,\bar{c}_{HW} \\ \bar{a}_{W\partial W} &= -2\,\frac{m_h^2}{2m_W^2}\left(\bar{c}_W + \bar{c}_{HW}\right)\end{aligned} \tag{E.135}$$

$$\begin{aligned}\bar{a}_{ZZ} &= -2\,\frac{m_h^2}{m_Z^2}\left(\bar{c}_{HW} + \bar{c}_{HB}\tan^2\theta_W - 4\bar{c}_\gamma \tan^2\theta_W \sin^2\theta_W\right) \\ \bar{a}_{Z\partial Z} &= -2\,\frac{m_h^2}{2m_Z^2}\left(\bar{c}_W + \bar{c}_{HW} + (\bar{c}_B + \bar{c}_{HB})\tan^2\theta_W\right)\end{aligned} \tag{E.136}$$

The amplitude (E.133) includes terms of $O(v^2/f^2)$, $O(m_h^2/M^2)$ and $O(\alpha_2/4\pi)$, while it neglects contributions of $O(v^4/f^4)$ and $O[(\alpha_2/4\pi)^2]$. As explained in Section 4.2, the corrections of $O[(\alpha_2/4\pi)(v^2/f^2)]$ are only partly included through the terms proportional to $\bar{c}_{HW}$ and $\bar{c}_{HB}$.

E.6 Numerical formulas for the decay rates in the SILH Lagrangian

We display here numerically approximated formulas of the Higgs decay rates valid at linear order in the effective coefficients $\bar{c}_i$ of the SILH Lagrangian (2.2) for $m_h = 125\,\mathrm{GeV}$. Although these numerical results have been partly shown in Section 3, we think it is useful to collect them here in a systematic and complete way. All the ratios Γ/Γ_{SM} have been computed by switching off the EW corrections, since their effect on the numerical prefactor appearing in front of the coefficients $\bar{c}_i$ is of order $(v^2/f^2)(\alpha_2/4\pi)$ and thus beyond the accuracy of the formulas implemented in `eHDECAY`. Conversely, we have fully included the QCD corrections, as they multiply both the SM and the NP terms. The numerical results are thus the following:

$$\frac{\Gamma(\bar{\psi}\psi)}{\Gamma(\bar{\psi}\psi)_{SM}} \simeq 1 - \bar{c}_H - 2\,\bar{c}_\psi\,, \tag{E.137}$$

$$\frac{\Gamma(h \to W^{(*)}W^*)}{\Gamma(h \to W^{(*)}W^*)_{SM}} \simeq 1 - \bar{c}_H + 2.2\,\bar{c}_W + 3.7\,\bar{c}_{HW}\,, \tag{E.138}$$

$$\begin{aligned}\frac{\Gamma(h \to Z^{(*)}Z^*)}{\Gamma(h \to Z^{(*)}Z^*)_{SM}} \simeq{}& 1 - \bar{c}_H + 2.0\,\left(\bar{c}_W + \tan^2\theta_W\,\bar{c}_B\right)\\ &+ 3.0\,\left(\bar{c}_{HW} + \tan^2\theta_W\,\bar{c}_{HB}\right) - 0.26\,\bar{c}_\gamma\,,\end{aligned} \tag{E.139}$$

$$\begin{aligned}\frac{\Gamma(h \to Z\gamma)}{\Gamma(h \to Z\gamma)_{SM}} \simeq{}& 1 - \bar{c}_H + 0.12\,\bar{c}_t - 5\cdot 10^{-4}\,\bar{c}_c - 0.003\,\bar{c}_b - 9\cdot 10^{-5}\,\bar{c}_\tau\\ &+ 4.2\,\bar{c}_W + 0.19\left(\bar{c}_{HW} - \bar{c}_{HB} + 8\,\bar{c}_\gamma\sin^2\theta_W\right)\frac{4\pi}{\sqrt{\alpha_2\alpha_{em}}}\,,\end{aligned} \tag{E.140}$$

$$\begin{aligned}\frac{\Gamma(h \to \gamma\gamma)}{\Gamma(h \to \gamma\gamma)_{SM}} \simeq{}& 1 - \bar{c}_H + 0.54\,\bar{c}_t - 0.003\,\bar{c}_c - 0.007\,\bar{c}_b - 0.007\,\bar{c}_\tau\\ &+ 5.04\,\bar{c}_W - 0.54\,\bar{c}_\gamma\,\frac{4\pi}{\alpha_{em}}\,,\end{aligned} \tag{E.141}$$

$$\frac{\Gamma(h \to gg)}{\Gamma(h \to gg)_{SM}} \simeq 1 - \bar{c}_H - 2.12\,\bar{c}_t + 0.024\,\bar{c}_c + 0.1\,\bar{c}_b + 22.2\,\bar{c}_g\,\frac{4\pi}{\alpha_2}\,. \tag{E.142}$$

F How to run eHDECAY: Input/Output Files

The program `eHDECAY` is self-contained, like the original code `HDECAY` on which it is based. All the new features related to the Lagrangian parametrizations proposed in this paper are encoded in the main source file, ehdecay.f, while other linked routines are taken over from the original version. Of course `eHDECAY`, besides calculating Higgs branching ratios and decay widths according to the non-linear, SILH or MCHM4/5 Lagrangian, also calculates the SM and MSSM ones, exactly as `HDECAY` 5.10 does. The choice can be done through the flags HIGGS and COUPVAR set in the input file. The input file for `eHDECAY` has been called ehdecay.in and is based on the file hdecay.in of the official version 5.10, supplemented by further input values. Thus, with the flag LAGPARAM the user can choose between the general SILH parametrization Eq. (2.2), the model-specific parametrizations MCHM4 and MCHM5 and the general non-linear Lagrangian parametrization Eq. (3.46). Furthermore, the various related couplings can be set. The input values are explained in the following:

COUPVAR, HIGGS: If HIGGS=0 and COUPVAR=1, then the Higgs decay widths and branching ratios are calculated within the parametrization chosen by:

LAGPARAM:

0: Non-linear Lagrangian parametrization Eq. (3.46)
1: SILH parametrization Eq. (2.2)
2: MCHM4/5 parametrization (cf. Table 1)

IELW: Turn off (0) or on (1) the electroweak corrections for the SILH, MCHM4 or MCHM5 parametrization. [30]

For the non-linear Lagrangian the following parameters have to be set for the couplings of the various vertices:[31]

---

[30] Note, that this parameter IELW has nothing to do with the parameter ELWK in the input file of HDECAY, where the meaning of this flag is different.

[31] We explain them here all, although they are in part already present in the input file for `HDECAY` 5.10.

CV: hVV vertex, (V=W, Z)
Ctau: $h\tau\tau$ vertex
Cmu: $h\mu\mu$ vertex
Ct: $ht\bar{t}$ vertex
Cb: $hb\bar{b}$ vertex
Cc: $hc\bar{c}$ vertex
Cs: $hs\bar{s}$ vertex
Cgaga: coupling $c_{\gamma\gamma}$
Cgg: coupling c_{gg}
CZga: coupling $c_{Z\gamma}$
CWW: coupling c_{WW}
CZZ: coupling c_{ZZ}
CWdW: coupling $c_{W\partial W}$
CZdZ: coupling $c_{Z\partial Z}$

In case of the SILH parametrization the input values to be set in order to calculate the various couplings are:

CHbar: $\bar{c}_H$
Ctaubar: $\bar{c}_\tau$
Cmubar: $\bar{c}_\mu$
Ctbar: $\bar{c}_t$
Cbbar: $\bar{c}_b$
Ccbar: $\bar{c}_c$
Csbar: $\bar{c}_s$
CWbar: $\bar{c}_W$
CBbar: $\bar{c}_B$
CHWbar: $\bar{c}_{HW}$
CHBbar: $\bar{c}_{HB}$
Cgambar: $\bar{c}_\gamma$
Cgbar: $\bar{c}_g$

In the MCHM4/5 parametrization we have the input values:

FERMREPR:
1: MCHM4
2: MCHM5

XI: the value for ξ

For example:

```
COUPVAR = 1
HIGGS = 0
⋮
************** LAGRANGIAN 0 - chiral 1 - SILH 2 - MCHM4/5 **************
LAGPARAM = 0
**** Turn off (0) or on (1) the elw corrections for LAGPARAM = 1 or 2 ****
```

```
IELW = 1
****************** VARIATION OF HIGGS COUPLINGS************************
CW = 1.D0
CZ = 1.D0
Ctau = 0.95D0
Cmu = 0.95D0
Ct = 0.95D0
Cb = 0.95D0
Cc = 0.95D0
Cs = 0.95D0
Cgaga = 0.005D0
Cgg = 0.001D0
CZga = 0.D0
CWW = 0.D0
CZZ = 0.D0
CWdW = 0.D0
CZdZ = 0.D0
⋮
```

computes the branching ratios for $c_V = 1$, $c_\psi = 0.95$ ($\psi = t, b, c, s, \tau, \mu$), $c_{\gamma\gamma} = 0.005$, $c_{gg} = 0.001$ and $c_{Z\gamma} = c_{WW} = c_{ZZ} = c_{W\partial W} = c_{Z\partial Z} = 0$ in the general parametrization Eq. (3.46). The output is written into the files br.eff1 and br.eff2, where the Higgs mass, branching ratios and total width are reported. For the previous example, at $m_h = 125$ GeV and for all the other parameters set at their standard values, the output reads

```
   MHSM        BB       TAU TAU     MU MU         SS          CC          TT
_______________________________________________________________________________

 125.000     0.5895     0.5654E-01 0.2002E-03 0.2161E-03 0.2569E-01  0.000

   MHSM          GG     GAM GAM     Z GAM         WW          ZZ         WIDTH
_______________________________________________________________________________
```

```
 125.000     0.9611E-01 0.1932E-03 0.1526E-02   0.2045   0.2554E-01 0.4129E-02
```

All the input parameters of the corresponding run are printed out in the file br.input. Otherwise, setting COUPVAR=0, the program produces the usual output files with SM or MSSM results according to the `HDECAY` 5.10 version.

References

[1] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716** (2012) 1 [arXiv:1207.7214 [hep-ex]].

[2] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716** (2012) 30 [arXiv:1207.7235 [hep-ex]].

[3] S. Weinberg, “The quantum theory of fields.” Cambridge, UK: Univ. Pr. (1996) 489 p

[4] G. F. Giudice, C. Grojean, A. Pomarol and R. Rattazzi, JHEP **0706** (2007) 045 [hep-ph/0703164].

[5] D. B. Kaplan and H. Georgi, Phys. Lett. B **136** (1984) 183. S. Dimopoulos and J. Preskill, Nucl. Phys. B **199**, 206 (1982). T. Banks, Nucl. Phys. B **243**, 125 (1984). D. B. Kaplan, H. Georgi and S. Dimopoulos, Phys. Lett. B **136**, 187 (1984). H. Georgi, D. B. Kaplan and P. Galison, Phys. Lett. B **143**, 152 (1984). H. Georgi and D. B. Kaplan, Phys. Lett. B **145**, 216 (1984). M. J. Dugan, H. Georgi and D. B. Kaplan, Nucl. Phys. B **254**, 299 (1985).

[6] R. Contino, Y. Nomura and A. Pomarol, Nucl. Phys. B **671** (2003) 148 [hep-ph/0306259].

[7] K. Agashe, R. Contino and A. Pomarol, Nucl. Phys. B **719** (2005) 165 [hep-ph/0412089].

[8] C. J. C. Burges and H. J. Schnitzer, Nucl. Phys. B **228** (1983) 464; C. N. Leung, S. T. Love and S. Rao, Z. Phys. C **31** (1986) 433; W. Buchmuller and D. Wyler, Nucl. Phys. B **268** (1986) 621.

[9] R. Rattazzi, Z. Phys. C **40** (1988) 605; B. Grzadkowski, Z. Hioki, K. Ohkuma and J. Wudka, Nucl. Phys. B **689** (2004) 108 [hep-ph/0310159]; P. J. Fox, Z. Ligeti, M. Papucci, G. Perez and M. D. Schwartz, Phys. Rev. D **78** (2008) 054008 [arXiv:0704.1482 [hep-ph]]; J. A. Aguilar-Saavedra, Nucl. Phys. B **812** (2009) 181 [arXiv:0811.3842 [hep-ph]]; J. A. Aguilar-Saavedra, Nucl. Phys. B **821** (2009) 215 [arXiv:0904.2387 [hep-ph]].

[10] C. Grojean, W. Skiba and J. Terning, Phys. Rev. D **73** (2006) 075008 [hep-ph/0602154].

[11] B. Grzadkowski, M. Iskrzynski, M. Misiak and J. Rosiek, JHEP **1010** (2010) 085 [arXiv:1008.4884 [hep-ph]].

[12] R. Barbieri, A. Pomarol, R. Rattazzi and A. Strumia, Nucl. Phys. B **703** (2004) 127 [hep-ph/0405040].

[13] R. Barbieri, CERN-TH-6659-92.

[14] M. E. Peskin and T. Takeuchi, Phys. Rev. D **46** (1992) 381.

[15] K. Agashe and R. Contino, Phys. Rev. D **80** (2009) 075016 [arXiv:0906.1542 [hep-ph]].

[16] G. Isidori, arXiv:1302.0661 [hep-ph].

[17] G. Isidori, Y. Nir and G. Perez, Ann. Rev. Nucl. Part. Sci. **60** (2010) 355 [arXiv:1002.0900 [hep-ph]].

[18] P. Sikivie, L. Susskind, M. B. Voloshin and V. I. Zakharov, Nucl. Phys. B **173** (1980) 189.

[19] G. Altarelli and R. Barbieri, Phys. Lett. B **253**, 161 (1991); G. Altarelli, R. Barbieri and S. Jadach, Nucl. Phys. B **369**, 3 (1992) [Erratum-ibid. B **376**, 444 (1992)].

[20] M. Baak, M. Goebel, J. Haller, A. Hoecker, D. Kennedy, R. Kogler, K. Moenig and M. Schott *et al.*, Eur. Phys. J. C **72** (2012) 2205 [arXiv:1209.2716 [hep-ph]].

[21] M. Redi and A. Weiler, JHEP **1111** (2011) 108 [arXiv:1106.6357 [hep-ph]].

[22] N. Vignaroli, Phys. Rev. D **86** (2012) 115011 [arXiv:1204.0478 [hep-ph]].

[23] M. Pospelov and A. Ritz, Annals Phys. **318** (2005) 119 [hep-ph/0504231].

[24] P. Paradisi and D. M. Straub, Phys. Lett. B **684** (2010) 147 [arXiv:0906.4551 [hep-ph]].

[25] J. F. Kamenik, M. Papucci and A. Weiler, Phys. Rev. D **85** (2012) 071501 [arXiv:1107.3143 [hep-ph]].

[26] J. A. Aguilar-Saavedra, N. F. Castro and A. Onofre, Phys. Rev. D **83** (2011) 117301 [arXiv:1105.0117 [hep-ph]].

[27] A. Hoecker and W.J. Marciano, PDG review on "The Muon Anomalous Magnetic Moment", in: J. Beringer *et al.* [Particle Data Group Collaboration], Phys. Rev. D **86**, 010001 (2012).

[28] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, Eur. Phys. J. C **71** (2011) 1515 [Erratum-ibid. C **72** (2012) 1874] [arXiv:1010.4180 [hep-ph]].

[29] D. Hanneke, S. F. Hoogerheide and G. Gabrielse, arXiv:1009.4831 [physics.atom-ph].

[30] G. F. Giudice, P. Paradisi and M. Passera, JHEP **1211** (2012) 113 [arXiv:1208.6583 [hep-ph]].

[31] G. W. Bennett *et al.* [Muon (g-2) Collaboration], Phys. Rev. D **80** (2009) 052008 [arXiv:0811.1207 [hep-ex]].

[32] J. J. Hudson, D. M. Kara, I. J. Smallman, B. E. Sauer, M. R. Tarbutt and E. A. Hinds, Nature **473** (2011) 493. D. M. Kara, I. J. Smallman, J. J. Hudson, B. E. Sauer, M. R. Tarbutt and E. A. Hinds, New J. Phys. **14** (2012) 103051 [arXiv:1208.4507 [physics.atom-ph]].

[33] R. Rattazzi, talk at the workshop *Physics at LHC: from Experiment to Theory*, Princeton University, March 21-24 2007

[34] R. Contino, C. Grojean, M. Moretti, F. Piccinini and R. Rattazzi, JHEP **1005** (2010) 089 [arXiv:1002.1011 [hep-ph]].

[35] R. Contino, L. Da Rold and A. Pomarol, Phys. Rev. D **75** (2007) 055014 [hep-ph/0612048].

[36] S. Y. Choi, D. J. Miller, 2, M. M. Muhlleitner and P. M. Zerwas, Phys. Lett. B **553** (2003) 61 [hep-ph/0210077].

[37] A. De Rujula, J. Lykken, M. Pierini, C. Rogan and M. Spiropulu, Phys. Rev. D **82** (2010) 013003 [arXiv:1001.5300 [hep-ph]].

[38] S. Bolognesi, Y. Gao, A. V. Gritsan, K. Melnikov, M. Schulze, N. V. Tran and A. Whitbeck, Phys. Rev. D **86** (2012) 095031 [arXiv:1208.4018 [hep-ph]].

[39] A. Azatov, A. Falkowski, C. Grojean and E. Kuflik, work in progress.

[40] J. R. Ellis, M. K. Gaillard and D. V. Nanopoulos, Nucl. Phys. B **106** (1976) 292; M. A. Shifman, A. I. Vainshtein, M. B. Voloshin and V. I. Zakharov, Sov. J. Nucl. Phys. **30** (1979) 711 [Yad. Fiz. **30** (1979) 1368].

[41] B. A. Kniehl and M. Spira, Z. Phys. C **69** (1995) 77 [hep-ph/9505225].

[42] A. Falkowski, Phys. Rev. D **77** (2008) 055018 [arXiv:0711.0828 [hep-ph]].

[43] I. Low and A. Vichi, Phys. Rev. D **84** (2011) 045019 [arXiv:1010.2753 [hep-ph]].

[44] A. Azatov and J. Galloway, Phys. Rev. D **85** (2012) 055013 [arXiv:1110.5646 [hep-ph]].

[45] M. Gillioz, R. Grober, C. Grojean, M. Muhlleitner and E. Salvioni, JHEP **1210** (2012) 004 [arXiv:1206.7120 [hep-ph]].

[46] M. Redi, Eur. Phys. J. C **72** (2012) 2030 [arXiv:1203.4220 [hep-ph]].

[47] R. Barbieri, D. Buttazzo, F. Sala, D. M. Straub and A. Tesi, arXiv:1211.5085 [hep-ph].

[48] C. Degrande, J.–M. Gerard, C. Grojean, F. Maltoni and G. Servant, JHEP **1103** (2011) 125 [arXiv:1010.6304 [hep-ph]]; H. Hesari and M. M. Najafabadi, arXiv:1207.0339 [hep-ph]; C. Englert, A. Freitas, M. Spira and P. M. Zerwas, arXiv:1210.2570 [hep-ph].

[49] C. Degrande, J. M. Gerard, C. Grojean, F. Maltoni and G. Servant, JHEP **1207** (2012) 036 [arXiv:1205.1065 [hep-ph]].

[50] S. R. Coleman, J. Wess and B. Zumino, Phys. Rev. **177** (1969) 2239; C. G. Callan, Jr., S. R. Coleman, J. Wess and B. Zumino, Phys. Rev. **177** (1969) 2247.

[51] C. P. Burgess and D. London, hep-ph/9203215.

[52] A. Azatov, R. Contino and J. Galloway, JHEP **1204** (2012) 127 [arXiv:1202.3415 [hep-ph]].

[53] R. Alonso, M. B. Gavela, L. Merlo, S. Rigolin and J. Yepes, arXiv:1212.3305 [hep-ph].

[54] R. Contino, C. Grojean, D. Pappadopulo, R. Rattazzi, A. Thamm, work in progress.

[55] G. Passarino, Nucl. Phys. B **868** (2013) 416 [arXiv:1209.5538 [hep-ph]].

[56] A. Djouadi, J. Kalinowski and M. Spira, Comput. Phys. Commun. **108** (1998) 56 [hep-ph/9704448]. A. Djouadi, M. M. Muhlleitner and M. Spira, Acta Phys. Polon. B **38** (2007) 635 [hep-ph/0609292].

[57] S. Weinberg, Physica A **96** (1979) 327.

[58] C. Grojean, E. E. Jenkins, A. V. Manohar and M. Trott, arXiv:1301.2588 [hep-ph].

[59] J. Elias-Miro, J. R. Espinosa, E. Masso and A. Pomarol, arXiv:1302.5661 [hep-ph].

[60] R. Barbieri, B. Bellazzini, V. S. Rychkov and A. Varagnolo, Phys. Rev. D **76** (2007) 115008 [arXiv:0706.0432 [hep-ph]].

[61] A. Pomarol and J. Serra, Phys. Rev. D **78** (2008) 074026 [arXiv:0806.3247 [hep-ph]].

[62] S. L. Adler, J. C. Collins and A. Duncan, Phys. Rev. D **15** (1977) 1712.

[63] J. C. Collins, A. Duncan and S. D. Joglekar, Phys. Rev. D **16** (1977) 438.

[64] N. K. Nielsen, Nucl. Phys. B **120** (1977) 212.

[65] M. Farina, C. Grojean, F. Maltoni, E. Salvioni and A. Thamm, arXiv:1211.3736 [hep-ph].

[66] K. G. Chetyrkin, B. A. Kniehl and M. Steinhauser, Nucl. Phys. B **510** (1998) 61 [hep-ph/9708255].

[67] M. Kramer, E. Laenen and M. Spira, Nucl. Phys. B **511** (1998) 523 [hep-ph/9611272].

[68] Y. Schroder and M. Steinhauser, JHEP **0601** (2006) 051 [hep-ph/0512058].

[69] K. G. Chetyrkin, J. H. Kuhn and C. Sturm, Nucl. Phys. B **744** (2006) 121 [hep-ph/0512060].

[70] T. Inami, T. Kubota and Y. Okada, Z. Phys. C **18** (1983) 69; A. Djouadi, M. Spira and P. M. Zerwas, Phys. Lett. B **264** (1991) 440; K. G. Chetyrkin, B. A. Kniehl and M. Steinhauser, Phys. Rev. Lett. **79** (1997) 353 [hep-ph/9705240].

[71] M. Spira, A. Djouadi, D. Graudenz and P. M. Zerwas, Nucl. Phys. B **453** (1995) 17 [hep-ph/9504378].

[72] P. A. Baikov and K. G. Chetyrkin, Phys. Rev. Lett. **97** (2006) 061803 [hep-ph/0604194].

[73] J. F. Gunion and H. E. Haber, Phys. Rev. D **67** (2003) 075019 [hep-ph/0207010].

[74] L. Randall, JHEP **0802** (2008) 084 [arXiv:0711.4360 [hep-ph]].

[75] J. F. Gunion and H. E. Haber, Nucl. Phys. B **272** (1986) 1 [Erratum-ibid. B **402** (1993) 567].

[76] K. Blum, R. T. D'Agnolo and J. Fan, JHEP **1301** (2013) 057 [arXiv:1206.5303 [hep-ph]].

[77] A. Azatov and J. Galloway, Int. J. Mod. Phys. A **28** (2013) 1330004 [arXiv:1212.1380 [hep-ph]].

[78] T. Appelquist and C. W. Bernard, Phys. Rev. D **22** (1980) 200; A. C. Longhitano, Phys. Rev. D **22** (1980) 1166; A. C. Longhitano, Nucl. Phys. B **188** (1981) 118.

[79] C. A. Baker, D. D. Doyle, P. Geltenbort, K. Green, M. G. D. van der Grinten, P. G. Harris, P. Iaydjiev and S. N. Ivanov *et al.*, Phys. Rev. Lett. **97** (2006) 131801 [hep-ex/0602020].

[80] W. C. Griffith, M. D. Swallows, T. H. Loftus, M. V. Romalis, B. R. Heckel and E. N. Fortson, Phys. Rev. Lett. **102** (2009) 101601.

[81] A. Sirlin, Phys. Rev. D **22** (1980) 971.

[82] E. Braaten and J. P. Leveille, Phys. Rev. D **22** (1980) 715; N. Sakai, Phys. Rev. D **22** (1980) 2220; T. Inami and T. Kubota, Nucl. Phys. B **179** (1981) 171; M. Drees and K.–I. Hikasa, Phys. Rev. D **41** (1990) 1547; M. Drees and K.–I. Hikasa, Phys. Lett. B **240** (1990) 455 [Erratum-ibid. B **262** (1991) 497].

[83] S. G. Gorishnii, A. L. Kataev, S. A. Larin and L. R. Surguladze, Mod. Phys. Lett. A **5** (1990) 2703; S. G. Gorishnii, A. L. Kataev, S. A. Larin and L. R. Surguladze, Phys. Rev. D **43** (1991) 1633; A. L. Kataev and V. T. Kim, Mod. Phys. Lett. A **9** (1994) 1309; S. G. Gorishnii, A. L. Kataev and S. A. Larin, Sov. J. Nucl. Phys. **40** (1984) 329 [Yad. Fiz. **40** (1984) 517]; L. R. Surguladze, Phys. Lett. B **341** (1994) 60 [hep-ph/9405325]; S. A. Larin, T. van Ritbergen and J. A. M. Vermaseren, Phys. Lett. B **362** (1995) 134 [hep-ph/9506465]; K. G. Chetyrkin and A. Kwiatkowski, Nucl. Phys. B **461** (1996) 3 [hep-ph/9505358].

[84] K. G. Chetyrkin, Phys. Lett. B **390** (1997) 309 [hep-ph/9608318].

[85] P. A. Baikov, K. G. Chetyrkin and J. H. Kuhn, Phys. Rev. Lett. **96** (2006) 012003 [hep-ph/0511063].

[86] J. Fleischer and F. Jegerlehner, Phys. Rev. D **23** (1981) 2001; D. Y. Bardin, B. M. Vilensky and P. K. Khristova, Sov. J. Nucl. Phys. **53** (1991) 152 [Yad. Fiz. **53** (1991) 240]; A. Dabelstein and W. Hollik, Z. Phys. C **53** (1992) 507; B. A. Kniehl, Nucl. Phys. B **376** (1992) 3.

[87] U. Aglietti, R. Bonciani, G. Degrassi and A. Vicini, Phys. Lett. B **595** (2004) 432 [hep-ph/0404071]; U. Aglietti, R. Bonciani, G. Degrassi and A. Vicini, Phys. Lett. B **600**

(2004) 57 [hep-ph/0407162]; S. Actis, G. Passarino, C. Sturm and S. Uccirati, Nucl. Phys. B **811** (2009) 182 [arXiv:0809.3667 [hep-ph]].

[88] G. Degrassi and F. Maltoni, Phys. Lett. B **600** (2004) 255 [hep-ph/0407249]; S. Actis, G. Passarino, C. Sturm and S. Uccirati, Phys. Lett. B **670** (2008) 12 [arXiv:0809.1301 [hep-ph]];

[89] H.–Q. Zheng and D.–D. Wu, Phys. Rev. D **42** (1990) 3760; A. Djouadi, M. Spira, J. J. van der Bij and P. M. Zerwas, Phys. Lett. B **257** (1991) 187; S. Dawson and R. P. Kauffman, Phys. Rev. D **47** (1993) 1264; A. Djouadi, M. Spira and P. M. Zerwas, Phys. Lett. B **311** (1993) 255 [hep-ph/9305335]; K. Melnikov and O. I. Yakovlev, Phys. Lett. B **312** (1993) 179 [hep-ph/9302281]; M. Inoue, R. Najima, T. Oka and J. Saito, Mod. Phys. Lett. A **9** (1994) 1189; J. Fleischer, O. V. Tarasov and V. O. Tarasov, Phys. Lett. B **584** (2004) 294 [hep-ph/0401090].

[90] G. Degrassi and F. Maltoni, Nucl. Phys. B **724** (2005) 183 [hep-ph/0504137].

[91] M. Spira, A. Djouadi and P. M. Zerwas, Phys. Lett. B **276** (1992) 350.

[92] R. N. Cahn, Rept. Prog. Phys. **52** (1989) 389.

[93] A. Bredenstein, A. Denner, S. Dittmaier and M. M. Weber, Phys. Rev. D **74** (2006) 013004 [hep-ph/0604011]; A. Bredenstein, A. Denner, S. Dittmaier and M. M. Weber, JHEP **0702** (2007) 080 [hep-ph/0611234].